

111

111

Maurilii Antonii
Bruno
Medicinae Doctor
1800.

VERMORE D. 46

VERRE E. 45

~~1003~~

N° d'inventario

371

371

X-K-22



U. S. Department of the Interior
Bureau of Land Management
Washington, D. C.

For the purpose of the survey of the
land in the State of California, the
following is a list of the names of the
persons who have been granted the right
to locate the same.

1. John A. Smith
2. James B. Jones
3. William C. Brown

4. Robert D. White
5. Thomas E. Black
6. Charles F. Green

7. Henry G. Hall
8. George H. King
9. Frederick I. Lee

10. John J. Scott
11. William K. Adams
12. James L. Baker

13. Charles M. Campbell
14. Thomas N. Clark
15. George O. Davis

16. Frederick P. Edwards
17. John Q. Foster
18. William R. Gibson

19. James S. Hall
20. Charles T. Jones
21. George U. King

22. Frederick V. Lee
23. John W. Scott
24. William X. Adams

25. James Y. Baker
26. Charles Z. Campbell
27. Thomas A. Clark

28. George B. Davis
29. Frederick C. Edwards
30. John D. Foster

31. William E. Gibson
32. James F. Hall
33. Charles G. Jones

*Stirpes, quas in Valle Stura Cuneensi,
Thermis Vinadiensibus, locisque Circumvicinis
Oviri tradidit Cl. Florae ped. Auctor Enumerantur;
Ordine ipsius Auctoris, servato.*

1. *Valeriana angustifolia.*
Coppione Cirenthermas Vinadii in sacris frequent in Valle Ma-
crae et Sturae. Incolit loca lapidea et meridiana sole percutta.
2. ——— *montana.*
In alpiibus Vinadiensibus
3. *Galium boreale.*
G. foliis quaternis lanceolatis trinerviis glabris caule erecto semine hispido.
Circa thelmas Vinadii in sacris Sylvatim.
4. ——— *rotundifolium*
prope thelmas Vinadii Cl. Bellardi.
5. *Salvia verticillata.*
Circa thelmas Vinadii.
6. *Thymus alpinus.*
Circa thelmas Vinadii frequent est.
7. *Satureja montana.*
In Valle Sturae frequent est.
8. *Lavandula spica*
ad Montes subalpinos Vinadii.
9. *Laminium ovata*
Circa thelmas Vinadii.
10. *Betonica Monierii*
In alpiibus Vinadiensibus.
11. *Melissa grandiflora.*
Circa thelmas Vinadii.
12. *Buglossum Barrelieri.*
In Valle Sturae prope Demont.
13. *Spinguicula lusitanica*
Sp. nectaris apice incrassato
Ex alpiibus Vinadiensibus Cl. Giavelli.
14. ——— *Villosa*
Sp. Scapo villosa flore minore calneo
Ex alpiibus Vinadiensibus Cl. Giavelli.
15. *Toxaria Alpina.*
Circa Vinadii thelmas in montibus le Vallon di S. Anna.

16. *Melampyrum Sylvaticum*
In alpiibus Vinadiensibus.

17. *Euphrasia officinalis*
C. L. alpina flava luteis floribus
In alpiibus circa Vinadii thermas

18. ——— *tricuspidata*
C. fol. linear. tricuspidatis
In alpiibus Vinadiensibus Cl. Gravelli.

19. *Pedicularis tuberosa*
P. caule simpliciter fl. spic. rostratis calyc. glabris aphodeti radice.
In alpiibus Vinadiensibus prope lacellum J. Annae.

20. *Scrophularia Vernalis*
In montibus Vinadii prope balnea.

21. *Linus alpinus* Calyc. 5-phyllus corol. pene infundibuliform. 5-fida vix
bilabiata segmentis emarginatis capsula 4-valvis polysperma
Didynamia angiospermia.
Circum thermas Vinadii.

22. ~~*Primula*~~ *Primula viscosa*
P. fol. viscosis odoratis linguiformibus sinuatis integerrimis
folia gemmae clementia palmatis subtiliter villosa Cynaglor. Vulgari
fol. similia Scapus palmatis flores coruleo pallidi 6. aut 7.
Calyc. communiter bidentem parvis foliis quot sunt radii
fructus maturus eximendus.
Hae Thyrse per errorem hic enumerata non ad Vinadii sed ad
Valderi thermas pertinet.

22. *Primula integrifolia*
P. fol. integerr. glabr. oblong. calycibus obtusis
In alpiibus Vinadii.

23. *Gentiana punctata*
G. coroll. subsidis campanul. punctat. Calycibus 5-dentatis
pallidum florem habet punctis distinctum fol. ovat. lanceol. nervos.
Circum Vinadii thermas.

24. ——— *palustris*
In Spongiarum lac de la Maddalene in extrema Vinadii Valle
Cl. Gravelli.

25. *Campanula latifolia*
Circum Vinadii thermas.

26. *Phytolacca Halleri*
P. foliis radicalibus cordatis acutis incisis tubis longissimis
hirsutis emarginatis spica ovata obesa fl. purpureo coruleo
Circum thermas Vinadii in pratis umbrosis.

Valle

Valle

Valle

Valle

Valle

Valle

Valle

27 Scabiosa alp
J. fol.
prop.

28 Cirsium
J. fol.
Alun.

29 —
J. fol.
Alun.

30 —
C. fol.
Alun.

31 Centaurea
C. cal.
J. fol.
Alun.

32 Artemisia
A. fol.
Alun.

33 Achillea
J. fol.

34 Chamæm
C. fol.
A. fol.
Alun.

35 Helian
J. fol.

36 Aster
J. fol.

37 —
J. fol.
Alun.

27. *Scabiosa alpina*
f. foliis pinnatis pinnis serratis capitulis globosis hirsutis floribus cernuis
prope Demonte loco dicti le Courmale

28. *Cirsium pyrenaicum*
f. foliis decurrentibus oblongo lanceolatis margine ciliatis et spi-
nosius utrinque tomentosis floribus tuberculatis.
Abunde circa thermales Vinadii.

29. — *helenoides*
fol. ciliat. subtus toment. radical. petiol. ovatis lanceolatis aut semi-
pinnatis
Abunde circa thermales Vinadii.

30. — *heterophyllum*
f. fol. amplexicaul. lanceol. ciliat. integris laciniatis. caul. subuni-
fol. inermi.
In jugo Col de la Madalene.

31. *Centaurea rhaphanifolia*
f. calycibus scariosis fol. ovatis-oblong. denticul. integris petiolatis sub-
tus tomentosis
Supra thermales Vinadienses editiora alpina parva Verius sanctuaria
f. Annae inhabitat.

32. *Artemisia vulgaris*
f. fol. pinnatis caul. adnudentibus fl. globosis cernuis recept. pappo.
Abrupta lammata alpium incolit et frequens est in Valle Vinadii.

33. *Achillea herbacea*
In montibus Vinadii.

34. *Chamaemelum montanum*
f. foliis late pinnatis pinnis palmatis caule folioso unifloro Hall.
f. fol. pinnatis-multifidis plan. lacin. linear. acut. hisp. pedunculo
longissimo.
Abunde in alpiibus Vinadii.

35. *Aster amellus*
In loco Vinadii dicto Nivajsa.

36. *Aster montanus*
In alpiibus Vinadiensibus

37. — *odoratus*
f. fol. amplexicaul. dentat. hirsutis. radical. ovatis. caul. lanceolatis
caule paucifloro. radice odora.
Circum thermales Vinadii.

38. *Hieracium aurantiacum* H. fol. integr. caule tubundo simpliciss. ppter. longum.
Circa thesma Vinadii. 6. fer.

39. *Scorzonera laciniata*
S. fol. linear. dentat. acut. caul. erect. squam. calyc. spatulo-erectis.
In montibus di. Demonte.

40. *Lunaria rediviva*
L. foliis alternis siliquis longioribus ovatis lanceolatis oblongis
Circa thesma Vinadii.

41. *Blatta hincoria*
Circa thesma Vinadii: ad rupes prope fontes thesmarum.

42. *Cardamine alavisolia*
Vinadiensis: alpis: familiaris.

43. *Sinapis maritima*
Circa thesma Vinadii:
fol. radical. long. petiolat. caulina parvis dentata sine triangulatis
summa hastata succosa crassula glaberrima caul. glaberrimus
calyc. spatul.

44. ——— *recurvata.*
S. fol. radicalibus subuncinatis caulinis pinnatifidis siliquis re-
curvatis
Ad thesma Vinadii: locis sterilibus.

45. *Lythrum hieracifolium* fol. lanceolatis serratis
In montibus Vinadiensis: frequens locis apertis

46. *Sisymbrium tanacetifolium*
In pascuis. umbratis circa thesma Vinadii: abunde.

47. *Epilobium dyneri*
Jecus aquar. ad thesma Vinadiensis.

48. *Anthyllis montana*
Ad rupes Berzel in Valle Vinadii.

49. *Ononis columnea*
O. fl. sessilibus fol. omn. ternat. stipul. lanceol. dentatis.
Vinadii e viciis Sarcorum La Brece

50. *Hedysarum alpinum*
Inter Vinay & le plance.

Page

100

101

102

103

Orabast
O. fol. p.
Citra

Hippoc
Zu

Latepikun
L. fol. p.
Zu v.

Liquicun
Zu p.

—
Zu

Jugonum
Abun

—
Abun

Myrrh.
Zu p.

Gerania
Citra the
5 p. 6 p.

Manne
Ac. fol.
teger
Zu

Stenhill
e. Scy

—
Zu

51. *Orobanchus luteus*
O. fol. pinnat. ovat. oblong. stipul. rotundato lunatis, caule simplici
Circa thesma Vinadii Cl. Gravelli.
52. *Hippocrepis uniliquata*
In apertis secus Ichiator supra Vinadii thesma
53. *Lathyrus gallicus*
L. foliis cuneiformibus furcatis.
In Valle fl. Thura prope Demont.
54. *Ligusticum meum*
In pratis alpinis Vinadiensibus
55. — *ferulaeum*
In alpidibus Vinadiensibus loco dicto le Barricade.
56. *Imyrenium olusatrum*
Abunde circa thesma Vinadii.
57. — *nodiflorum*
Abunde in sylvis proximisque pratis Vinadii.
58. *Myrrhis odorata*
In pinguibus pratis alpium Vinadii.
59. *Geranium sylvaticum*
Circa thesma Vinadiense in Valle di Lemie.
G. ped. biflor. fol. subserrat. 5. lob. inci. serrat. caul. erect. petal. emargin.
60. *Ranunculus thora*
R. fol. inferior. subrotund. reniform. crenat. floral. ~~exsert.~~ lanceolat. integerrim. caul. infera. nud. caul. subflor.
In alpidibus Vinadiensibus.
61. *Potentilla Canadensis*
e scopulis Vinadiensibus prope la Madonna del Vallon praevalit.
62. — *Valdetia*
In saxosis alpium Vinadii.

63. *Potentilla grandiflora*
Circum Thermas Vinadii.

64. *Delphinium elatum*
Circum Thermas Vinadienses in umbratis locis non rarum.

65. *Aquilegia viscosa*
In alpinis Vinadii: Cl. Bellardi.
A. caule subnudo subunifloro foliis ternatis foliolis subtilibus sessilibus.

66. *Helleborus foetidus*
In alta Valle Thurae ruscensis.

67. *Saxifraga Cotyledon*
S. folior. ora cartilaginea caule triplicato variegato petalis immaculatis
In Valle Thurae loco dicto le Batticade.

68. *Dianthus superbus*
Loco dicto Col d'Y mul. D. Giavelli.

69. *Silene cordifolia*
Circum Thermas Vinadii

70. *Viola alpina*
Folia dentata uniformia ovata hanc ab aliis affinis distingunt
Cl. Bellardi: Secus torrentem Eschiator prope Thermas Vinadii legit.

71. *Cistus Serpyllifolius*
C. suffruticosus stipulatus fol. herbac. venos. marginibus parvis ciliatis
Circum Thermas Vinadii in rupibus

72. *Dictamnus albus* Decandria Monogynia L.
Cal. 5. phyllis deciduis corollae 5. petala subpapilionacea l. petal. descendente Capsulae 5. margin. inter. Coarctae superius deliscentes
Covinatae polyspermae semina nigra Calyptra elastica involuta.
D. fol. pinn. caul. simplicis: Odor naturalis vultus communis
Petala magna patentissima ex albo purpurascens.
In alpinis Vinadiensibus Cl. Giavelli.

73. *Sedum rupestre*
Circum Vinadii Thermas locis siccis et apricis
S. foliis subulatis 5. fariis confectis basi solutis floribus cymosis.

74. *Ribes alpinum*
In dumetis et montanis proximis pratis, circa Vinadii Thermas.

len. lib.

calabi

ingruat
legit

calabi

et al de
centu
muni

fil.

mat

75. Ribes gl.
A. var.
Grossul.
Ab.

76. Allium
A. sea
Bn.

77. Tulipa
T. fl.
Bn.
Hgh.

78. Salix m.
J. fo.
Bn.

79. Alnus
Circ.

80. Rosa L.
R. sp.
Bn.

81. Festuca
F. rac.
muc.
C. A.

82. Pteris
F. fo.
legu.
Bn.

83. Polyp.
P. sp.
Bn.

84. Ulva
U. c.
Bn.

85. Cacti.
Lo.

75. *Ribes grossularia*

A. ramis aculeatis, petiolorum ciliis pilosis, bacis hirsutis
Grossularia fructu minimo hispido margaritarum fere colore.
 Abunde circa Vinadii Thermas.

76. *Allium triquetrum*

A. scapellum fol. triquetris stamin. simpl. caule triangulari.
 In alpiibus Vinadii Cl. Gravetti.

77. *Tulipa sylvestris*. Corolla campanulata petal. oblongo cuspidatis, concavis.

T. flor. subnata fol. lanceolatis

In pratibus alpibus Vinadiensibus.

Stigma triangulare angulis divitis.

78. *Salix myrsinites*

L. foliis serratis glabris ovato lanceolatis venis julis tomentosis

In alpiibus Vinadii.

79. *Alnus europaea*

Circa Thermas Vinadienses.

80. *Poa sellesoides*.

sp. spica ovata locustis subserilibus sepalis obtusis.

In pratibus elatioribus Vinadii Cl. Bellardi.

81. *Festuca phaeocoides*

F. racemo indiviso spicul. altera. n. brevilibus teretibus foliis involutis
 mucronato pungentibus

Cl. Bellardi in Valle Sturae prope Dement.

82. *Pteris Crispa*

F. fol. duplicato pinnatis sterilibus obscure dentatis, floralibus ovatis in-
 tegerrimis.

In montibus Vinadii frequentissima.

83. *Polypodium Pontederæ*

sp. pinnis paucis semipinnatis brevissimis acute serratis

In Sylvis umbrosis ad Thermas Vinadii

84. *Uva labyrinthiformis* fructificationes habet in membrana diaphana

U. cellulis labyrinthiformibus prominentibusque clavatis.

In thetunis Vinadii grad. lat. 50.

In Auctario Cl. Adlonius nullis plantis hæc natali
 loco assignat Thermas Vinadienses nec Vallam / Stura

In Appendice

85. *Cacalia albifrons*

Locis humidioribus Vinadii.

Cunei 1800
 Die 26. Maji.

Addenda

86. *Ranunculus palustris*
in foliis circa Demont.
planta graminifolia lamina aquatica:

87. *Cyananthus alpinus*
singularum varietatem vel distinctam speciem reulit D. Balbise
propositum per alpes Vinadii in hibernia, quod videtur
Leucanthemum pyrenaicum minimum multifido incano folio.

77

lie.

Betonica Monieri hirtula L.

B. spica majore hirtior labio superiore emarginato plano stamina saepe
longiuscula Chaise

B. spica bari foliola, corollis galea integra L.
obli. floribus purpureis differt a B. alopecuroides L. quod flores habet luteos.

Veronica austriaca

Caulis ferre perispermatus villis folia varie ludunt plerumque
abrotani similia quandoque Coronopifolia amulantur glabra racemi
longi Calyces 5fid: Corolla pallide vel saepe cerulea

Phytneuma Haller

Rapunculus foliis radicalibus cordiformibus acute incisis tubis longis
simis hirtis emarginatis spica ovata obesa. Haller.

Artemisia rupestris

A. foliis pinnatis caulibus adscendentibus floribus globosis (ceruis)
receptaculo papposo Linn.

Abrutivum etc alio florum foliis angustis sericeis pinnatis pinnis
3. fidis & 5. fidis Haller

Artemisia mutellina foliis petiolatis multiplicis floribus pedunculatis
axillaribus terminalibusq. receptaculis hirtis calyce anguloso villis.

Enumeratio plantarum, quas in Valle Gessii,
prope Valdevii thermas, Oppida Rutagne, Roascia,
et Vicina loca crescere adnotavit Cl. Allionius.

1. *Lavandula spica.*
ad montes subalpinos Valdevii
2. *Lomium ovula*
Circum thermas Valdevii in saxosis umbratis.
3. *Betonica menierii*
In alpi. bus Valdevianis
4. *Melissa grandiflora*
Circum thermas Valdevii copiosissime.
5. *Melampyrum Sylvaticum*
in summa sylva umbrosa thermis tepidi Valdevianis imminente
abunde.
6. *Veronica austriaca* ~~ll~~
V. racem. lateral. fol. linear-lanceol. pinnat. dent. tenuiter. laciu.
Cl. Bellardi in Montibus Roascia
Mac V. verna varietas est All. auch.
7. *Junculentra*
V. erecta spicata, fol. succul. caulibus pinnato-dent. floral. integerrim.
In saxosis appicis circum thermas Valdevianas
8. *Samolus Valevandi.* Pentandria Monogynia.
Calyx 5 fidus Corolla 5-fida rotata squama muniens 5. Capsula
unilocularis infera polysperma. 5 Valvis.
Caul. ped- cubital. etiam ramulos fulco virens fol. petiol. obovata
tuberculata nitida fl. racemosi albi Caps. fulca glabra.
Circum fontes calidiores Valdevii.
9. *Primula Vircola.* Vide plantas Vinadienses.
Circum thermas Valdevii ad rupes.
10. *Gentiana asclepiadea*
Circum Valdevii thermas
11. *Campanula ulicifolia*
in umbratis sylvis Circum Valdevii thermas
12. *Phitheimma Halleri*
Circum thermas Valdevii in pratis umbratis
13. *Artemisia vulgaris* abrupta summorum alpinum incolit &
est frequens in valle Valdevii. & Vinadii

14. *Achillea herbacea*
In montibus Valdevii.

15. *Chamaemelum montanum*
In alpiibus Valdevii.

16. *Matricaria inodora*
M. receptac. hamisphaer. radii patentibus seminibus coronato marg.
ginalis squamis calycinis margine obsolete.
Differt. a M. chamomilla habitu firmiori: defectu odori
majori flore
In montibus Valdevii.

17. *Linaria halmifolia* Vide *Silpes Valli Maiva*.
Alyrium fol. lanceol-lineur. acut. integerr. caul. procumb.
perennant. flores racemosi primum umbellati albidissimi.
Siliqua ovata compressa.
In Montibus di. Roasca Cl. Bellardi.

18. *Cardamine alarifolia*
familiavit alpiibus Valdevianis.

19. *Sisymbrium tanacetifolium*
Circa theumas Valdevii: Copiose.

20. *Brachiolobos pyrenaicus*
in alpiibus Valdevii.

21. *Smignium nodiflorum*
abunde in sylvis procerisque pratis Valdevii.

22. *Potentilla valderia*
frequent in saxosis alpiibus Valdevii.

23. *Saxifraga stellata*
abunde circa theumas Valdevianas.

24. *Saxifraga pedemontana*
Hujus insignis Varietatis naccibus circa Theumas Valdevii
foliis 5. lobis, lobo medio saepe 2. aut 3. pido petalis albis ad
ungues roseis

25. *Silene cordifolia*
Circa theumas Valdevii.

Chamaemelum montanum

C. foliis laevibus pinnatis pinnis palmatis caule folioso (-floro. Haller.
Anthemis foliis pinnato-multifidis planis laciniis linearibus acutis
bispidis pedunculo longissimo Lin.

Caul. palm-terquiped. ramolus incurvus Folia Carrociniana apice
latiora, & tantum a medio ad apicem pinnata laciniis ovatis acutis non
linearibus. Ad basin petiolorum pinnata bipartita ut in A. arvensi.
Flores Chrysanthemi Leucanthemi Vitman.

Saxifraga pedemontana All.

S. caule erecto subnudo foliis trilobis & bidentatis calicinis palmatis
Ad radicem haec habet rotundam foliorum quae ex angustis principio
triangulariter amplificantur & in tres lineares lobos profunde secantur & si
lobi bidentati sunt raro 4. dentati. Caulis inter ea erigitur folio uno circa
medium donatus quod manus instar in 4. Lobos dividitur umbellatim
quasi flores ex summo caule nascuntur ad pedunculorum orthum quin-
que digito aut tridigito foliolo opposito. Sed singuliflores tridentur
pedunculo ter floribus longiore, & nudo. Flor. albus spatens petalis ex
linguae ovatis etiam ellipticis calyce duplo longioribus. Calyx lobatus
profunde sectus in laciniis lanceolatas. Stamina cum antheris luteo
virentibus calyce non altiora. Tota planta viscida & hirsuta est
Circa Thermas Valderii eadem planta aliam faciem induit
atque folia minime gerit profunde bipida sed pinnata 5-loba
loto medio saepe bispido aut 3-fido atque petala alba ad
linguae rostra sunt Allion

Viola nummularifolia
V. foliis cordato-acuminatis glabris integerrimis.
V. fol. subrotund. obiter crenatis stipulis lanceolatis serratis caule bre.
vicinis bifloro.

Obi. Folia in hac viola sunt cordato acuminata glabra integerrima
 flori violæ marisæ æqualis stipulæ tridentata. Allion.

Cette plante est très-petite elle n'a que 2. ou 3. pouces de haut
 & les pedoncules sont les deux tiers de sa longueur, la racine est
 vivace noirâtre & noueuse les feuilles sont presque rondes avec
 une pointe presque insensible & une échancrure encore plus
 petite à leur base elles sont toutes pétiolées n'ont que 3. à 4.
 lignes de diamètre & autant de longueur elles ont à leur base
 2. stipules lanceolés & dentés à dents aiguës & sensibles qui
 regardent le sommet la fleur est petite penchée, bleu lavé ou
 blanchâtre, le nectar est obfus la moitié plus long que le calice

Willard

Viola Valdenia

V. erecta herbacea foliis ellipticis sinuatis non dentatis stipulis inte-
gerrimis

Radix hujus violæ est parva & albicans etc quæ cauliculus erectus vix unum
 vnum duarum altitudinem habens striatus & angulosus. Folia alternatim &
 dense nascuntur. Radicalia aut inferiora pene orbiculata sunt reliqua ellip-
 tica, aut longe elliptica petiolo pene longitudine folii & inferius sulcata.
 Foliorum aversa facies venosa est ambitus autem sinuatus & integerrimus Flores
 etc alii foliorum nascuntur pedunculis plantæ altitudinem superantibus fere bi-
 uncialibus Flor paullo minor mihi videtur et quam in *V. Calcarata*. Tubus ap-
 crassus est & perforatus & stipulæ præstant duo foliola sæpius inæqualia.
 Obsecro habet folia sinuata non dentata stipulæ etiam integras.

Allion

26. *Viola nummularifolia*
Inter diffractos lapides summi et ediculi jugi Col de la femamorta
ad alpes Valderianas pertinentis enascitur. Locis summi & alpe-
ribus quos pertinent ad alpes d'entraive lantolla Tenda
& Madonna delle Finestre.

27. *Viola Valdevia*
Circa Valdevii thermas in Saxosis torrentis ad radices montis. Boaniti.
Copiose circa la Madonna delle Finestre.

28. *Stellaria hypericifolia* Vide *Stellaria Valdevia*
Copiosissime circa thermas Valdevii.

29. *Sedum alpinifolium* Vide *Stirpes in Valle Macra valcentis*
Cl. Bellardi: in montibus di Roalchia.

30. *Rubus cassinus*
Circa thermas Valdevii.

31. *Tulipa sylvestris*
in pratis alpini Valdeviani.

32. *Asarum europaeum*
Circa thermas Valdevianas.

33. *Pteris Crispa*
in montibus Valdevii frequentissima.

34. *Polypodium dryopteris*
filiis ramolaminos. pinnulis dentatis
frequentissima circa thermas Valdevii.

35. *Uva labyrinthiformis*
in thermis Valdevii gradu Caloris 50.

Auctarii

36. *Buglossum Barvetieri*
in Vallibus di Vaudier et Entraque

37. *Phytolacca Carnetii* Vill.
prope Entraque in regione dicta la Lanza.

38. *Hieracium ruppelii*
in montibus d'entraive locis Saxosis et Calidis dictis la Lanza
H. fol. ovatis. pilos. dentatis. laciniatis. caul. sub 1. floro non plio
calycibus subincaut.

39. *Hieracium Anthybaceum*
Circa thermas Valdevianas.

40. *Turris stricta*
T. glabra lucens caule simplicissimo erecto siliquis planis margine
attenuatis
prope thesma Valderii

41. *Selinum Seguietii* L.
S. involucri universi nullo spiritibus diuaticatissimi
prope Entragne.

42. *Geranium phaeum* prope Entragne.

43. *Cucubalus alpestris*
In alpestribus di Entragne e rupium fissuris prodit locis umbris
et daucos

44. *Viola nummularia* folia
locis humilis et alpestribus quae pertinent ad Alpes di Entragne.

45. *Epipactis atrorubens*
L. caule aphyllis flore demissis labello ovato lanceolato calcar ovato
turgido.
labium baccis compressis dent. caul. vagin. corollae resupinatis ne-
ctarii labio indebito
In fagibus prope Entragne locis umbris sub humido.

46. *Corallorhiza neottia*. Genus Valde affine *Epipactis*: differt an-
theris ymatensis
Ophrys bulb. vacuol. flosculis. caul. vagin. aphyll. nectar. lab. 3. fil.
obovato radice coralloide vuberrima.
In pratens molineti in fagibus prope Entragne.

47. *Zunera filiformis*
locis paludosis di Entragne.

48. *Festuca altissima*.
F. fol. plan. villosis albo nervo divinis; panicula contracta pedunc.
communibus geminatis baccis trifloris.
locis lapideis & umbris prope thesma Valderii et in Valle di
Entragne prope Locellum S. Joannis.

Cucubalus Alpestri? All.

C. dichotomus foliis linearibus calycibus pelvisimilibus inflatis striatis
petalis bipartitis.

Radice parum fibrosa inter rumpum suam oblique repit perennis atque
plures caules prostratis qui primum geniculati sunt deinde se erigunt
ad palmarem altitudinem rotundi glabri simplices: tota etiam planta glaber-
rime est nihil habent viscosi aut villosi. Folia linearia acuminata
glabra mollia ad 2. fere pollices longa de more gentis opposita & con-
nata caulis summus praebet perennium aut alteram dichotomum
flores singulares pedunculatos: pedunculus lateralis est brevior non bra-
cteatus: calyce ovato five peltatis pelvisimilibus sursum ampliores inflatis
glabri altera parte rubentes ore breviter 5-dentato dentibus ophelii.
petala esse solent introflecta profundissime bifida ex his sulco rubentia
intus sordide alba. Nylites squamula nulla coronatice sed eorum loco
(non quidem semper duo parvi tumores.

Exhibet habitum suo *J. Alpestrem* Jacq. sed haec & squamula habet et
petala bifida calycis etiam diversitatem. *Alliden*

Festuca altissima

Radix biennis & ultra. Radicis primariae truncus plures teretes fibras demittit. Ge-
summa radice plures culmi teretes prodeunt ad 3. & ultra pedum altitudinem.
los vertunt undique vagina foliorum usque ad prostratum folium. Haec
aspera sunt & asperitudine sua superant foliorum superficiem & ambitum
culmorum vaginis denudatis laevit & striatissima est. 4. circiter folia in unoquoque
culmo esse solent. Sed praeter haec ad culmi basim 2. folia adiuva erecta
multum reliquis breviora. Folia caulina recta ex vagina discedunt fere
horizontalia 3. lineas & ultra latitudinem. Subtus nervo albo divisa plana in
aristam denique acuminata. Ad latera culmorum solet oriri fasciculus
foliorum altero anno novam plantam daturus quae pedalis & ultra al-
titudinis sunt Culmi sunt simplices & ferunt paniculam non patentem
sed collectam pedunculis erectis. pro planta altitudine brevis panicula
est quae scilicet 4. uncias fere non superet. Pedunculi primi

¶ secundari geminati nascuntur secundi pedunculi alternatim pro-
ferunt locustas tres aut quatuor, quarum prima pedunculata est alij
sessilibus. Haec locustae interdum biflorae saepe 3-florae glabrae acumi-
natae. Calyx flore brevior & angustior est glumis angustis acuminatis
persistenteribus, altera gluma multum brevior. Corolla florum eie glum-
is aequalibus glabris extensori sub aristata facile caduca semen
gracile — Allion Auctar

9. Lichen
f. l.
July

49. Lichen Cornucopioides.
L. Scyphifer simplex folio brevior tuberculis coccineis
Zutylui prope thesaur. Valderii locis umbratis et rupibus.

Cunei 27. Maji.
1800.



Handwritten text at the top of the page, possibly a title or header, reading "H. ... in Halle ...".



Itinerario in Valle d'Aosta

[Faint, illegible handwritten text, likely bleed-through from the reverse side of the page.]

Observationes.

Valeriana monandra Vill.

Valeriana angustifolia Les feuilles sont linéaires, très étroites, souvent par paquets avec aisselles des autres feuilles. Elle vient dans les endroits pierreux, sur les graviers, le long des torrents, des montagnes. Villars.

Lunaria biarmifolia

Cespitem facit cauliculorum vis erectorum robustior. Semipalmatis aut etiam aliorum. folia longe elliptica crassula concava glandulosa granulata cinereo virentia sessilia flores racemosi primum umbelati fl. albissimi patentissimi petal. 7 ovatis sep. levill. emarginatis. Siliqua ovata compressa semina marginata. — All.

Scorzonera luciniata

l. fol. linear. dentatis pinnatifidis acutis caule erecto squamis calycinis patulo mucronatis
La tige est perpendiculaire droite & ramifiée Les fleurs sont jaunes petites dans un calyce argutense dont les écailles sont terminées par un bouton cotonneux. Elle vient dans les lieux escarpés exposés au midi & paron: les Alpes. Villars

Levisimum hiernifolium L.

La tige est haute de huit ponce environ simple & non ramifiée Les feuilles inférieures sont dentées un peu blanchâtres ou cendrées celles de la tige deviennent linéaires. Les fleurs d'un jaune pâle terminent la plante. Les siliques sont dures un peu velues & rapprochées de la tige. Elle vient dans les montagnes pierreuses Villars

*Stirpes quae in Valle Maerae
Crescere videntur. Cl. Allionius.*

1. *Valeriana angustifolia*
V. fol. integr. longissim. et angustiss. flor. calcar. denato.
In Valle Maerae.
2. *Satureja montana*
In Valle Maerae legit Terraneus.
3. *Primula longiflora* / Similis *primula farinosa*.
sp. fol. glabr. elliptic. rugos. serrat. flor. umbell. tubo longissim.
In Valle Maerae prope la Brusà & in alpidi Vraita.
Segmenta floris habet purpureo-violacea - A. A.
4. *Erica herbacea*
Frequent in Valle Maerae.
5. *Atropa belladonna*.
In Valle Maerae et lot.
6. *Santolina chamaecyparissus*
S. pedunculis unifloris, foliis quadrifariis dentatis.
Franciscus Peyroletti in Valle Maerae Collegit.
7. *Senecio parviflorus*.
In alpidi Vallis Maerae.
8. *Cineraria alpina*
in Valle Maerae.
9. *Scrophularia laciniata*
In Valle Maerae locis siccis ad Viar prope Stopp.
10. *Lunaria halimifolia*
In aridis calidioribus locis Vallis Maerae.
11. *3beris Garreoliana*.
In Valle Maerae circa Stopp.
12. *Erysimum hieracifolium*.
In montibus Vallis Maerae frequenter locis apricis.
L. foliis lanceolatis serratis l.

13. *Trifolium pannonicum*.
In hum. dir. alpiibus di. Staj. et la Chianale.

14. *Ononis Cenisia*
In summa Valle Maera loco dicto la Ciappa frequens est.
O. glabra inermis, stipulis, dentato-vaginantibus, corollam vespillo reflexo
purpureo, alis & carina n. v. vill.

15. *Astragalus Uvalensis*
A. scapi, apophyllis, fol. ovato lanceolatis, sessilibus, siliquis hirsutis,
hirsutis erectis ~~styliferis~~ styliferis. Haller.
In alpiibus la Chianale.

16. *Pimpinella pumila* Genus involucri null. semina intus plana, paulisper
acuminata, dorso gibbo tricostata.
Frequens in Valle Maera
Tragoselinum caul. crass. sulcat. divaricato fol. multif. capill.
Sp. umbell. numerosissimis simplicissimis, compositisque

17. *Cichus pilosus*.
In Valle Maera nascitur.

18. *Sedum alpinefolium*. (a)
Circa S. Damian in Valle Maera.

19. *Acorus Calamus*. *Hexandria monogynia* L.
Gen. Clava ex sinu folii orivunda digitiformis, flor. sessil. un-
dique tecta calice 6phyllis, foliolis obtusis, superne crassioribus
persistentibus stylis brevissimis capsula triangulatis, trilocularis
seminibus 3. feta.

Locis spongiosis in Valle Maera lectus est a Peyroletti.
Radix horizontal. articulata. fol. radical. vaginantibus, ensiform. integris
odorata. Scapus apice foliosus. folii laevi sulcatum proferit spatheum
nudum digitiform. cylindric. subnudentem fl. in spica attenuata.
Variat clavis ternis, quaternis.

Cunei die 27. Maji 1800.

(a) *Sedum alpinefolium*
S. erectum ramosum foliis planis, floribus alb. longe pe-
dunculatis Allion.

Observationes

Trifolium pannonicum

T. spicis villis longis corollis monopetalis calyce erecto

villis villis juncy.

Cette espèce s'élève à un pied environ: les feuilles sont pointues très-velues ainsi que la tige: celle-ci se termine par un ou deux épis arrondis, que des corolles très-relevées blanches ou jaunâtres beaucoup plus grandes que dans les autres espèces font paraître allongés — Villars

Ononis laevis.

O. pedunculis unifloris multio foliis ternatis cuneatis ligulatis serratis caulibus prostratis

Cette espèce est petite quoique ligneuse elle est entièrement glabre excepté le calyce les folioles sont allongues avec des dentelures aiguës les fleurs naissent aussi à l'extrémité des feuilles supérieures: la corolle est glabre courte ventrée très-peu allongée. Villars

Asragalus luteus

As. acutis scapo erecto foliis longiore leguminibus subulatis inflatis villis erectis. Vill.

La racine est grosse mais simple ne produisant qu'une plante pour l'ordinaire les feuilles ne sont pas couchées par terre mais relevées le long du scape les scapes sont longs de six pouces & un peu plus droits & terminés par un épi de fleurs jaunâtres (ochroleucis). Les gousses sont droites rapprochées de la tige oblongues velues & pointues. Elle vient parmi les prés fertiles des Alpes dans les endroits herbueux & humides Vill.

Pimpinella pumila — *Dioica L.*

Caulis divaricatus palmatis laevissimus parvis foliatis folia ternata live bignata linearia. Umbellae terminales & laterales 2. oppositae plerumque compositae paucis simplicibus petala longe lanceolata incurva non emarginata Stamina alba longa involucrum nullum

Dans la plante mâle & hermaphrodite — G. libert.
M. Plant.

[Faint, mostly illegible handwritten text in a historical script, possibly Latin or Greek, covering the main body of the page. The text is organized into several horizontal sections separated by lines.]

[Marginal notes in a historical script, likely Latin, located on the right side of the page. The notes are written vertically and include botanical or geographical references.]

Qlen
et ad ja
hoccavi
Ede
dium boreale
prope Ten
Operula horap
et foliis deni
Ad vuppi
Thymus alpinu
Inter lino
— piperu
ad oppida
Sideritis Rou
herbacea
rari ovate.
Cl. Bellard
Laminum orsal
supra oppi
— gargar
okher cir
Bryolostum B
inter ho
Sclerophalaris
Circu limon
— Citrus alpi
Inter Te
Androsace v
in altitudinib
— Elica herba
in subalpi
— Alhiza bo
In Vall
Thymale
supra Te

Clenebus stirpium, quæ in Montibus Tenda,
et adjacentibus Territoriis Carlin, Briga, Limon,
Roccavione, Boves, Valperio &c. nascuntur.
Æthimonio Cl. Fl. Piedemontaneæ auctoris.

1. *Galium boreale*
prope Tenda in herbido ad Mineræ fodinam.

2. *Asperula hirsutifolia*
A-folii scilicet linearibus floribus umbellatis terminalibus subsessilibus
Ad vupes prope Tendam nascitur.

3. *Thymus alpinus*
Inter Limonem et Tenda copiose.

4. *Piperella*
ad oppida Carlin la Briga etc. e rupibus prodit.

5. *Sideritis Romana*
J. herbacea decumbens ebracteata calycibus spinosis; Labiis super-
riori ovato.
Cl. Bellardi in monte Tenda.

6. *Lamium orsala*
supra oppidum Limon

7. *Garganium*
oritur circa Limon.

8. *Buglossum Bartheletii*
inter Roccavione & vobitant.

9. *Scrophularia vernalis*
Circâ Limonem frequent

10. *Erinus alpinus* Vide Plant. Vinadii.
Inter Tenda et Savorgio.

11. *Androsace villosa*
in altioribus montibus supra Tenda.

12. *Erica herbacea*
in subalpinis sterilibus supra Tenda.

13. *Atropa belladonna*
In Valle di Perio prope monasterium charthusianorum.

14. *Thymus alpinus*
supra Tendam et Limonem ad vupes Ignatiæ et Molineri.

15. *Thymalæa cheorum*
In montibus di Tenda.
16. *Scabiosa alpina* Vid. Pl. Vindob.
Plurima provenit in Valle di Perio & montibus di Tenda e la Buga
17. — *graminifolia*
Erysimum fissurij prodit inter ulmeta et Carlino observ. Cl. Dana.
Supra Limone etiam occurrit.
18. *Globularia nudicaulis*
G. fol. oblonge ovatis integerrimis caule nudo unifloro
Cl. Bellardi in Montibus di Tenda.
19. *Lappula tomentosa*
Prope Tenda frequens est.
20. *Carduus acanthoides*
abunda prope Tenda.
C. alis caulinis latiss. fol. semipinnat. pinn. angulis spinosis flor.
longis fasciculatis
21. — *Collinoideus*
C. fol. decurrentibus decussatis pinnatis pinnulis palmatis 2-3
dentatis lanatis caul. corymb. multiflor. flor. glomeratis.
Circum Tenda
22. *Cirsium medium*.
Carduus fol. decurr. pinnatis marg. spinos. caul. 1-floro pedunculo
longill. calycem squamis internodibus patentibus setaceis.
In montibus di Tenda
23. — *Ochroleucum*.
C. fol. pinnatis pinnis ciliatis. Squam. calycin. recurvis.
Indescentia di Col di Tenda Cl. Bellardi.
24. — *canum*
Carduus fol. decurrent. lanceol. ex toto dentatis ciliatoaculeatis
utrinque arachnoideo-subvillosis.
in montibus Tenda.
25. — *helenioides*
C. fol. ciliatis subtus tomentosis radiculis petiolatis ovato lanceol.
aut semipinnatis
Ignobis Molinari circa Tenda.

2 labij

Dana

anij / lot

4 f. 20

/

20000

/

(2000)

anuel

cul / evor
foliis decur
Abunde in

laurea tha
supra find

temilia lo
lacus Vian

achillea he
nullibi

— m
supra J

typanthem
in montib

Mutricaria
Inter li

Cineraria
In mon

— la
Inter J

Arnica C
A. uniflo
Inter

Hieraciu
H. fol. inte
In me

Prenant
Gen. Cal
quot in
Vimplex
P. folio
Vimplex

Scoroner
In mon

Tragopo
T. calyco
Non in

26. *Cnicus fevoris*
C. foliis decurrentibus singulatis dentatis spinosis (a. b. l. e. d. a. m. o. s. e. r. e. t. o.)
Abunde in montibus circa Tenda.

27. *Centauria thapsus* V. J. Vinadianus.
Supra Tendam loco dicto ripreid.

28. *Alchemilla lobelia*
Juncus Viam inter Tendam & Nicam.

29. *Achillea herbacea*
nullibi copiosius quam supra Tenda.

30. ——— magna
Supra Tenda copiose nascitur.

31. *Gysanthemum Ceratophylloides*
in montibus Tenda et Briga

32. *Matricaria inodora* Vide plant. Valder.
Inter limen et Robitant.

33. *Cineraria alpina*
In montibus Tenda Briga, limone

34. ——— longifolia
Inter Tenda et la Madonna delle fenestre.

35. *Arnica Clusii*
A. uniflor. fol. altern. ov-lanceol. hirs. integer. calyc. hirs.
Inter Tenda & limone.

36. *Hieracium elvauliacum*
H. fol. integr. caul. subnud. simpliciss. pilos. corymbifero.
In montibus Tenda et laioribus.

37. *Prenanthes viminea*.
Gen. Calyce cylindraceo ex paucis squamis, fere parallelis cum ali-
quot minoribus ad basim. Semiflosculis in normis staminibus. Pappus
simplex stipitatus aut sessilis.
P. foliorum ramentis, caul. adnatis, flor. sessil. planta villosa
prope Tenda.

38. *Scorzonera laciniata* Vide pl. Vinad.
In montibus di Tenda

39. *Tragopogon exaristatus*
T. calycibus coroll. rad. longior. fol. integr. radical. perennantiq. bas. villor.
Non infrequens a jugo Col di Tenda ad Nicam.

40. *Bidentella auriculata*
B. Calycibus nectarario unilingue gibbis siliculis in stylum coeuntibus.
Circum Brigam Tendam.

41. *Thlaspi loricata*
Supra la Briga

42. *Blechnum garretianum*
Supra tendam in montibus editis inter Brigam & Callium.

43. *Cardamine alavisana*
Secum torrentes supra Tendam.

44. *Sinapis maritima* Vid. Plant. Vinad.
in decembris di. Col di Tenda.

45. — *Jounefortii*
Brassica fol. runcin. hispida. caul. hispida. siliq. torulosa. glabra. patent.
rostris longissimo.
Locis rupibus propeta Madonna delle fenestre.

46. *Heperis Sylvestris*
In sylvis caducis circa Limone

47. — *laciniata*
H. caul. ramosus. fol. inaeq. laciniatis. flor. sulphureo.

48. *Anthyllum montanum*
Circum Ceram supra Tendam

49. *Ononis villosa*
Abunde provenit Tendae locis aridis
O. pedunc. uniflor. aristat. fol. simpl. infim. ternatis.

50. *Cytisus lupinus*
in montibus ad Tendam pertinentibus
C. fl. umbellat. terminal. ram. decumbent. foliol. ovatis.
C. fl. axillar. ram. decumb. siliq. moll. lanug. pubescentibus.

51. *Genista pilosa*
Inter habitant et Roccamare
G. fol. lanceol. obtus. caul. tubercul. decumb. inermis.

52. *Asparagus depressus*
A. aculeos. scap. fol. brevior. legum. cernuis foliol. subemarginatis. nudis.
(C. Bellardi) Copiose crescere vidit in iugo Col di Tenda dicto.
Escalpin di Piero Petrus Cornalia.

14.

14.

14.

62 Peucedanum
p. fol. 4
In comitibus

63 p. fol. pin.
Jupia

64 Carassipitium
l. fol. lip.
inter res

65 Juniperium
ligusticum
verh. illa
Abund

66 Myrtus
Cider

67 Geranium
Jup

68 G. pedunc.

69 Malva m.
Inter

70 Ranunculus
Ranunculus
nodoso
ad petio
p. v. g. i.
In m.
p. de ca.

71 A. fol.
ceolato
Cl

72 Potentilla
p. fol.
Loric

53. *Pencedanum officinale*
p. fol. ~~tripartitis~~ quinquies 3 partitis filiformibus linearibus
In comitatu nicæensi supra Tenda.

54. ——— minus
p. fol. pinnat. foliol. pinnatifid. lacin. linear. oppos. caul. ramor. lim. patulo.
Supra Tenda.

55. *Lactucium gallicum*
l. foliol. cuneiformibus pinnatis
inter Tenda & Melicena

56. *Jungermannia nodiflorum*
ligusticum nodiflor. fol. cordat. profunde lobat. acut. caul. ramif.
verticillato umbell. inordinat. numerosiss.
Abunde Circa Tenda, Limone.

57. *Myrtus* ad ovata
Circa Limone

58. *Geranium macrorrhizon*
Supra Tendam observavit Ignatius Molineri.

59. ——— pyrenaicum
g. peduncul. 2. flor. fol. multifid. lacin. obtus. inaequal. pet. 2. fid.

60. *Malva moschata*
Inter Tenda & Limon.

61. *Hanunculus columba*
Hanunculus fol. orbiculatum multifid. caule prostrato ad genicula
nodoso fol. circumspersione robusta a flavo-ventria glauca splend.
ad petiol. usque 5. fida caul. breviss. brevis pedunculo 1. flor.
porrigit fl. lachent amplius antheras luteas.
In montibus di Limone loco dicti Vermelina ex diffraeto la.
pide Calcareo supra Castin. Cl. Dana.

62. ——— alpestris
A. fol. radic. sulcorat. obtus. 3. partitis lobis 3 lobatis caul. lan-
ceolato integerrimo caule subuni floro.
Cl. Dana in monte Relegà vestitus Tendam.

63. *Potentilla fruticosa*
p. fol. pinnat. caul. fruticosa.
Locis apricis in montibus l' Inferno dicti ad Tendam prestinentib.

64. *Potentilla caulescent.*
Supra Tendam occurrat.
65. *Potentilla Valdensis*
Supra Tendam
66. *Aquilegia Virgata*
In alpiniis Tendae.
67. *Jasione pedemontana*
J. caule erecto subnudo fol. tr. lob. et trident, caulibus palmatis
In editis jugis d. Tenda et limore.
68. *Silene Cordifolia*
Supra Tenda.
69. *Viola palmata*
V. acaulis fol. reniform. glabris.
Coppice ad finem scaberrime circa Tendam.
70. *Arenaria montana* idest. *Capillacea.*
A. fol. linear. lanceolat. scabr. caulibus steril. longis. prostratis.
Alpine fol. linear. acuminat. petal. flor. integr. calyc. duplo longioribus.
In humo jugo Tendae ha. sola dicta
71. ~~Linum~~ *Petraea*
A. fol. ovali? carinatis recurvis quadrifariam imbricatis. flor. aggregat.
In montibus Tendae locis siccis.
Fl. in capitul. congest. 5. vel 6.
72. *Sedum alpine folium.*
Sedum vameorum fol. plan. flor. alb. longe pedunculatis
Inter Tendam & Robilant.
73. *Allium arenarium*
A. caule planifolio bulbifero vagini; ferebunt spatula multa stami-
nibus bristlydatis
A. fol. ciliat. umbella bulbifera stamin. alterne bristly.
Secus torrentes proximamque prata circa Tendam.
74. *Allium nigrum*
in monte Vermelina prope Limon.
75. *Fritillaria meleagris*
in pratis alpiniis Tendae.
76. *Tulipa Sylvatica*
In pratis alpiniis Tendae.
77. *Semprevivum hirtum.*
S. fol. caule petalorumq. apicibus hirtis
Uberrime in montibus ad Tendam spectantibus.
78. *Buscus sempervirens*
inter Robilant & Rocavion.

Handwritten text in a cursive script, likely a letter or a page from a manuscript. The text is written in a dark ink on aged, slightly yellowed paper. The handwriting is fluid and characteristic of the 17th or 18th century.

Continuation of the handwritten text. The script remains consistent, with some variations in line spacing and ink density. The text appears to be a single, continuous piece of writing.

Further continuation of the handwritten text. The cursive script is well-maintained throughout the page, showing a high level of skill in the writer's hand.

The final section of handwritten text on the page. It concludes the piece with a few final lines of cursive script, followed by a small flourish or signature at the bottom.

[Faint handwritten text, likely bleed-through from the reverse side of the page. The text is mostly illegible due to fading and the angle of the page.]

[Handwritten notes in the right margin, continuing from the main text or as separate entries.]
 Scleroidy
 praelation
 cur melli
 glum. 2 flori
 hiculata.
 In palcat
 de Baum
 lypsa Cadu
 lve alere
 a. aphylla
 lrcul 2-3
 uli Capr. 3-
 In Cochit
 glottum Bo
 In Valle
 Antichinu
 A. fol. plevi
 inhabitat
 Nimula glub
 p. fol. Jov
 Cicala e
 Campanula
 Thymelaea
 in Vap: bar
 Dicoic
 Daphne flo
 Supra Teni
 Chamaemel
 Iningo ca
 Doronicum
 in montib
 Hieracium
 in prabi
 in
 Supra la
 Arabis as
 A. caule
 silig. fere
 ad oral

79. *Poa Selivoides* V. p. *Gradii*
in pratensationibus Tenda.

80. *Holcus mollis*
H. glum. 2 flori; nudiuscul. florum hermaphrodit. Anther. masculo arista
geniculata.
In pratens. elatioribus di Tenda Cl. Bellardi.

81. *Buxbaumia aphylla*. Gen. Anthera operculata hinc membranacea
Calyptra caduca inter operculum laevis pollinis. Capsul. Ovali
latere altero oblique plana orificio denticulato.
B. aphylla pedunculo elongato capitulo latio bivalvi.
Inscul 2-3 lineari. alter durius radice minimi ad basin tripartita.
talis cap. 3-lineari viridi.
In corticibus circa Tendam lecta est et D. Allionio communis.

in Auckavia notata.

82. *Buglossum Barrelieri*
In Valli: limone ubique.

83. *Antirrhinum origanifolium*
A. fol. pleriq. opposit. oblong. fl. v. altern.
inhabitat rupes meridie spectantes supra Tenda.

84. *Primula glutinosa*
p. fol. serrat. glutinos. calycum laciniis capsula longioribus.
Circa Madonna delle finestre.

85. *Campanula latifolia* in pinguis pratens. di Petio.

86. *Thymelaea lanamunda*
in rupibus meridie spectantibus supra Tenda.

87. ———— *dioica*
Daphne flor. axillar. geminis fol. linear-lanceol.
supra Tenda in rupibus meridie spectantibus.

88. *Chamaemelum montanum*
In jago Madonna delle finestre.

89. *Doronicum austriacum*
in montibus di Petio.

90. *Hieracium pratense*
in pratens. edioribus, della Madonna delle finestre

91. ———— *intybacum*
supra la Madonna delle finestre

92. *Arabis aspera*
A. caule erecto flexuoso ramoso fol. ovat. oppos. semiamplexicaulibus
siliq. teretibus.
Ad oral sylvarum caducarum supra Tenda.

93. *Arabis scabra*
In alpidibus di Tenda.
94. *Turritia stricta*
Supra Limone
95. *Vicia onobrychioides*
abunde locis montanis & alpidibus di Tenda atque montibus di Riveda.
96. *Bypleurum petraeum*
B. caule nudo fol. gramin. involucri et involucellis distinctis polyptichis.
In rupibus supra Limone loco dicto Armellin et circa la Madonna delle finestre.
97. *Chaerophyllum Sylvestris*
prope Castrum Spiriti.
98. *Geranium phaeum*
prope Robilant & Limone B. Molineri.
99. *Saxifraga mutata*
plurima est locis umbratis humidis rupibus supra la Madonna delle finestre.
100. *Dianthus glaucus*
in pascuis apricis et alpidibus di Tenda & Madonna delle finestre.
101. *Cucubalus alpestris*
C. dichotomus fol. linear. calyc. pelvitorm. inflat. striat. petal. bipartitis.
In alpidibus di Tenda & rupibus firmissimis prodit locis umbratis & siccis.
102. *Pyrola uniflora*
Supra Tenda.
103. *Viola nummularia folia*
locis summis alpidibus quae pertinent ad Alpes Tenda & Madonna delle finestre.
104. — *Valderia*
Coppione circa la Madonna delle finestre.
105. *Cirrus humulatus*
C. erichipulatus suffrutescens. flor. subumbel. petal. lunata crocea notat.
fol. oval.
In extrema Valle Spiriti, in montibus supra Limone.
106. *Fritillaria involucreata*
F. fol. opposit. summi. tenu. supra Tenda & il Riveda locis siccis
& rupibus mare spectantibus.



uncus filit
Loc. 7 pal

12. Fehaco heru
F. fol. rad
sub hirsuti
Loc. 11

1. Avena
~~A. p...~~
~~fol. A.~~
ti foliis
Abund

1. Avena
Avena lu
varicat
In pa

1. Hordeum
Circa Ca

2. Carex di
Zucolit
lem di

13. —
frequen
lexium

14. Lichen
L. umbril
Lapidib

15. Lichen
L. fruti
cautibus sc
Rupre
Donna de

116. Lycopu
hajus

117. Poa
p. pa
volu
Zuc

107. *Juncus filiformis*
locis paludosis alpinis di Tenda.

108. *Festuca heterophylla*
F. fol. radical. setaceis, caulinis planis, panicula erecta glumis
subhirsutis, margine scabris, subaristatis.
locis montanis Saceris montium di Tenda.

109. *Avena sempervirens*
~~A. paniculata patens locis paludosis alpinis di Tenda.~~
A. paniculata calycibus trifloris, hermaphroditis, bari-lana-
tis foliis viridibus acutis, involutis, sempervirentibus.
Abund. in montibus di Castell. la Briga.

110. *Avena bromoides*
Avena subspicata: spiculis binatis, altera pedunculata arctis di-
varicatis calycibus 8-floris.
In palceis Saceris inter il Rifredo & Valpelio.

111. *Hordeum europaeum*
Circa Carthesium di pelio locis sylvestribus B. Molineri.

112. *Carex dioica*
Incolit prata montana di Rifredo quae est Tenda itur in Val-
lem di pelio ad fontium frigidos & scaturigines.

113. — Bellardi
frequens locis alpestribus, solique patentibus alpinis Montega-
ledium versus Castell.

114. *Lichen polyphyllus*.
L. umbilicatus polyphyllus. undique laevi, atro-virens, venatus.
Lapidibus adhaerens, lectus est prope la Madonna delle Finestre.

115. *Lichen trisporus*
L. haticulatus solidus, ramulis compressis, ramulis apicibus nigri-
cantibus scutellis, terminalibus.
Rupes inhabitat, locis editis & vento perflatis, prope la Ma-
donna delle Finestre.

Plantae quae continentur in Appendice.

116. *Lycopus exaltatus*
hujus varietas incana nascitur serus v. vultus circa Provisum.

117. *Poa violacea*
P. panicula erecta stricta spiculis compressis, sub 4-floris co-
volvis, basi hirsutis.
In montibus limoni locis picis.

118. *Poa stolonifera*
P. panicula expansa spiculis compressis seris flori; culmo stolonifera
Nascitur in montibus Limoni et Vallispirii.
119. *Festuca flavescens*
Non in frequen; locis sylvaticis subalpini; Vallispirii Limoni
F. panicula secunda erecta flavescens spiculis oblongis acutis
in 65 floris fol. Capillat.
120. *Avena setacea*
A. fol. setae. panicula purpurascens calycibus trifloris
aviliis niguis recurvis.
In montibus Limoni locis calidioribus invenit D. Viale
121. *Galium pusillum*
in Valle spirii sed planta major.
122. ——— *montanum*
G. fol. tenuis lineari acuminatis glabris caule prostrato
pedunculis trichotomis floribus subumbellatis.
locis montanis Vallispirii Limoni.
123. *Asperula cyanantha*
stujus varietas vel distincta species crevit in rupibus
Vallispirii Limoni Tendae.
124. *Plantago argentea*
In montibus Vallispirii elatioribus parum a lanceolata diversa.
125. *Supplementum incurvum.*
D. Viale in montibus limonenis.
126. *Hyacinthus botryoides*
Fr. ligo prope Carthuriam.
127. *Juncifraga lingulata*
J. fol. radicalibus longissimis lineariibus sulcat. cartilagineis integris.
caule foliis paniculato.
In Valle spirii, & montibus Limoni.
128. *Juncifraga diaphanoides.*
In Valle spirii Fr. ligo in alpiibus limoni D. Viale. supra
Tendam de Juffren.
129. *Dianthus hirsutus*
D. flor. subaggregat. squam. calyc. vix aequant. lobis lobulatis
inversis caule foliisque laevibus.
D. caule fol. scabri flor. panicul. squam. avitatis calyce
brevioribus
In Monte Blimanda supra Bovitium locis sterilibus
130. *Aolapimpinellifolia*
A. glab. globos. glab. peduncul. par. rect. petal. scabr. foliis obtus
in montibus Tendae D. de Juffren.

intercept

Rubus glau
A. fol. spin
mirell
Zn mon
te Brill

Fr. lya

Anemo
Zn ad

Ranunc
A. fol.
temi ang
in pra
D. Vian

Gerani
Zn

Genit
G. fol. l
Zn l

Heij
Variet

Cacali
inter 2

Artem
prop

Erige
L. cau
in m

Ank
A. e
Non
Pili

Cen
C. ca
caul
Zn

Caren
Zn l
Non
C. lpi
mud

131. *Rubus glandulosus*
 A. fol. pinnat. ternatis simplicibusque caulium spinis rubris villosis
 mixtis fructibus cymosis hirsutis
 In montibus Vallis Pissii supra & circa Carthusiam. Copiose in mon-
 te Brimanda supra Bovitium.
132. — *triphyllos*
 Fr. ugo in Valle Pissii locis asperioribus.
133. *Anemone dubia*
 In alpidibus Limonensibus D. Viale
134. *Ranunculus lacerus*
 A. fol. radical. petiolat. cuneiform. laceris apice multifidis caulinis
 semiamplexicaulis laciniatis summit. semilibus linear. caul. laev. ramos.
 in pratis summorum alpidum Limonensium locis humidiusculis
 D. Viale.
135. *Geranium fuscum*
 In pratis Vallis Pissii.
136. *Genista scoparia*
 G. fol. simpl. hirsut. ram. 10. angulat. cinereis.
 In rupibus limoni a Sole perculis
137. *Hedysarum onobrichis*
 Varietas incana fol. longior nascitur in montibus Vallis Pissii
138. *Cacalia albifrons*
 inter diffractos lapides Vallis Pissii limoni
139. *Artemisia camphorata*
 prope Rocca di Vico.
140. *Erigeron Villarii*
 E. caule erecto fol. oblong. lanceolat. binerviis fl. corymbos.
 in montibus Carthusiae Vall. Pissii Fr. ugo. Vaso.
141. *Anthemis altissima*
 A. erecta fol. pinnat. pinnar. 6-8 denticul. reflexis asperis
 non infrequenter circa Bovitium inter segetes huminagris Vallis
 Pissii prope Carthusiam
142. *Centauria nudicaulis*
 C. calyc. setac. - spinos. squamos. fol. indivis. superior. subdent.
 caule nudiusculo simpliciter
 In montibus Vallis Pissii supra Carthusiam locis sylvest. Fr. ugo.
143. *Carex foliacea*
 In Valle Pissii prope Carthusiam locis humidiusculis D. Viale ex
 montibus limoni
 C. spicul. subovat. sessil. remot. aedrogyn. caput. ovah. ferebimus
 mutic. divaricat.

144. *Salix pontedera*.
Sal. fol. dur. oblong-lanceolat serrat. glandulos intern. glaucis, jatis
oblongis basi foliosis.
In altitudinis limoni D. Viale.
145. — *Daphnoides* !
f. fol. elliptico acuminatis, subserratis, inferne glaucis, superne nitidis
julis brevibus dense tomentosis.
In Valle Spisii et in montibus limoni. 147. Pteris Cretica
in Sylva cretica umbrat. Tendon.
146. *Polypodium limborum*
In monte Bissimanda supra Bovirium. in Valle Spisii Fr. Ugo.
147. *Picea minima* frondibus glabris bipartitis acutis
in Valle Spisii supra Carthusiam locis humidis.
148. *Lichen peltiphyllus*
f. subimbricat. foliol. orbiculat. lobatis. rub. marg. alb. tuberc. nigris.
In altitudinis Vallis Spisii Fr. Ugo.
149. — *Leucigerus*
f. crustaceus albicans sublobat. scutell. albo-margin. asate flavescens.
In Valle Spisii
150. — *Carpitum*
f. imbricat. cinereo-virescent. sublobato margine albo tuberculis cartil.
lenitissimis saepe congehit
Ex montibus limoni D. Viale
151. *Lichen velleiformis*
f. umbilicatus subtus hirsutus peltis imbricatis subtus saeculis
supra Jarcia in montibus limoni D. Viale.
152. *Boletus hirsutus*
Agaricum squamosum superne hirsutum & obscurum subtus etc. pulv.
aureum densissime et tenuissime perforatum.
In Valle Spisii ad caudices Castaneorum.
153. — *Cinnabarinus*
B. acutis pulvinatis tomentis et foris ruber
Fr. Ugo reperit supra Huncos emarcidos in Valle Spisii.
154. — *Medulla spani*
B. crustaceus effusus farinosus albus in montibus Bovirii & Valle Spisii.
155. *Clavaria cinerea* ramis rimosis apic. multis.
ad caudices arborum praescentium prope Bovirium.
156. *Lycoperdon epidendrum* cortice farinaceo purpureo
supra Huncos emarcidos fagi Sylva cretica in Sylva prope Ch. Vall. Spisii.
157. *Sphaeria fragipodii*
Ex montibus Carthusiae Fr. Ugo. plantarum. In adultis nigra.

Cuneidi 29. Maji 1800.



Cera.

1. *Calceola*
in *flor.*
in *flor.*

2. *Thymus*
in *flor.*
in *flor.*

3. *Thymus*
in *flor.*

4. *Thymus*
in *flor.*
in *flor.*
in *flor.*
in *flor.*
in *flor.*

5. *Lamium*
in *flor.*

6. *Teucrium*
in *flor.*

7. *Symphoricarpos*
in *flor.*

8. *Conium maculatum*
in *flor.*
in *flor.*
in *flor.*
in *flor.*

9. *Scrophularia*
in *flor.*

10. *Monarda*
in *flor.*
in *flor.*
in *flor.*
in *flor.*
in *flor.*

Plantae quae in Provincia Monregalensi et circa
Ceva, Garretto, Ormea recensere adnotavit Cl. Allionius
in Flora sua Pedemontana.

1. *Valeriana tuberosa*
N. foliis triandris; fol. radical. lanceolatis integerrimis; reliquis pinnatifidis.
Montes montana radice olivari.
Inter Garretto & Ormea loco dicti la fana dei Ovi.
2. *Thymus piperella*
In alpiibus montanis; quae pertinent ad Monregal. provinc. ad oppid.
Garretto.
3. *Hyssopus officinalis*
Oritur inter Garretto & Ormea.
4. *Thymus spicata* Gen. Cal. subcylindric. 2. labiat. utrinq. lineavit-
lola excavatus. Corol. galea obtusa 2-fida barba 3-fida, laciniis fere
aequalibus stylis semibifidis.
Cult. montibus inter Garretto & Ormea.
Caul. dodrant. lignori, gepiculati in can. Fol. hyssop. virid. atroviscent.
minoribus intermixtis foliol. in sum. Caulib. margin. laevigi-
natis. Fl. ex axillis bractear. tenui purpur.
5. *Laminum ovata*
in montibus umbratis Monregalensi provincia.
6. *Tenerium lucidum*
in sylvis Garretianis copiose.
7. *Symphytum tuberosum*
F. foliis summis oppositis
in montibus Monregalensibus copiosissime.
8. *Coris monspeliensis* Gen. Calyc. 10. dent. Coroll. parum fingen 5-fida
capitata. Valv. pentandria monogynia.
Circum Garretianum.
Caul. fruticosus palmis. ruber fol. altern. spars. linear. patentia
crassiuscula succulenta ciliata Fl. capitata spicati corollae ex
purpur-coerule.
9. *Scrophularia veruata*
copiose in montibus Monregalensibus
10. *Montia fontana* Triandria 3-gynia
Calyc. 2. 3-phyllus corollae 1. petal. irregul. parva. 5-part. lacin.
3. altern. min. Capsul. 1-locula 3. valv. sapient 3-lyserum.
ad fontium scaturigines circa Garretianum.
Caul. procumb. ramis. articul. in prat. viridis semiped. in gla-
veol. subune. Fol. petiol. ovata vel oblong. flor. racemos. pendul.
exiguus albi raro se se asperientes.

11. *Mimula longiflora*
In alpiibus Garrescio te Cavente
12. *Lysimachia nemorum*
ad Torrentem oppidi: Monastero Monti Regali
13. *Campanula bononiensis* — petra in alpiibus di Cusotto.
C. fol. ovato-lanceol. subtus scabris sessilibus caule paniculato.
In aridis locis Grappiol dicitur supra Garrescio.
14. *Phyteuma Halleri*
in stratis humosis alpinum Monregalensium
15. *Thymelea alpina*
Inter ormeam et Garrescium Zynaticum Molineri
Daphne flor. sessil. aggregat. lateral. foliis lanceolatis ob.
hiscensulis sub his timentolis.
16. *Poterium spinosum*
P. spinis ramosis
In montibus circa Garrescium non in frequent.
17. *Onopordum rotundi folium*
In alpiibus editionibus Monregalensibus
18. *Carduus acanthoides*
Circum Garrescio.
19. *Carduus acanthoides*
C. fol. decurrentibus decurvis pinnatis pinnulis gradatis 4. pidi
aculeatis lanatis. Cauli corymb. multiflor. floribus glomeratis.
Circum Garrescio.
20. *Cirsium Carniolicum* *Carduus rivularis* L.
Carduus fol. inferne pinnatis pidi in laciniis oblongis superne integris
& serratis inermi-spinulosis sessilibus caule inermi stipitibus.
ad fluentem aquam supra Garrescium.
21. *Artemisia lobelia*
Supra Garrescium loco dicto il Garbo della luna.
22. *Artemisia pontica*
Supra Garrescium cum precedenti eodem loco crevit.
23. *Chrysocoma linosyris* Gen. Calyce hemisphaerico imbricato. Fl. her.
maphrod. uniformes. 5. pidi Stylus vix florentis longior Receptac.
hancum seminum passim simplex.
C. herbacea fol. lineari. glabr. calyc. lac. squarrosis.
Planta digitata vix odorem aromaticum spirat.
In provincia Monti Regali saepe nascitur.

Ho.

206

11

reg

Fl. kn

hc.

[Faint, illegible handwritten text, likely bleed-through from the reverse side of the page.]

[Faint, illegible handwritten text, likely bleed-through from the reverse side of the page.]

24. *[illegible]*
25. *[illegible]*
26. *[illegible]*
27. *[illegible]*
28. *[illegible]*
29. *[illegible]*
30. *[illegible]*
31. *[illegible]*
32. *[illegible]*
33. *[illegible]*
34. *[illegible]*
35. *[illegible]*
36. *[illegible]*

24. *Xanthoxylum annuum*
In aridis montanis & collinis Monti Regalis sat frequens
25. *Matricaria parthenium*
In vineis Monti Regalis frequentissima.
26. *Aster officinalis*
In agro S. Albani Legit Cl. Bonelli.
27. — *montanus*
In alpebus Garreici, & territoris Monti Regalis.
28. — *odoratus*
Buxula fol. amplexicaud. dentat. hirsutis. radical. ovatis. caul. in lan-
ceolatis. caul. pauciflor.
Circa S. Michele Monti Regalis.
29. *Cineraria cordifolia*
In montibus Garreici Villar Souran dictis.
30. — *longifolia*
Supra Garreicio
31. *Epimedium alpinum*
Gen. electaria 4. cyathisperm. petalis incumbentibus Calyce caduco
4. phyllis corol. 4. petal. lappata acuminata 1. locut. 2. Valv.
poly 11.
In montibus Garreici
Caul. caudat. internod. juveniles diducit fuscus pallidus. Fol. long. petiol.
falsigata 3. ternata foliol. cordiform. dentat. Laxa virid. glabra
venosa rigida Fl. purpureo flavo variegati emarginati.
In montibus Garreici
32. *Draba nemorosa*
In montibus Monti Regalis frequens.
33. *Lunaria holymifolia*
Alyssum fol. lanceolat. linear. acut. integerr. caul. procumb. peren.
In collibus editioribus Garreici.
34. *Thlaspi alliaceum*
T. siliculis tuberculis ventricosis fol. oblong. obtus. dentat. glabr.
In vineis Circa Mondovi & Garreicio.
Planta allium redolens Caul. pedat. fol. infer. petiol. caul. in-
terit. vel amplexicaud. fl. fasciculati subumbellati longe pedun-
culati cor. alba.
35. *3beris garreiciana*
Supra Frabosa
36. — *jascaensis*
3. Jastrutaceus fol. lanceolat. linear. cavis, acut. integerr. ciliat.
Inter Jasca alpium Casotto Vicinarum.

37. *Cytisus lupinus*
 Super Ormea & Gavresio
 c. flor. umbell. peduncul. ram. decumb. foliol. ova.
38. *Genista pilosa*
 G. fol. lanceolata. obtus. cauf. tuberculat. decumb.
 loci aridi montium Monti Regali.
39. *Vicia Gerardii*
 In sylvestribus subalpini Gavresii dicti i pra di fior.
 V. pedunculat. multiflor. petiol. polyphyll. foliol. villos. stipul.
 acut. integ. appendiculat.
40. *Artragalus depressus*
 In alpiibus di Carso.
41. *Malva moschata*
 In apricis sylvi Gavresii et praesepue loco dicto il pra di
 fior.
42. *Fragaria sterilis*
 in montibus Monti Regali.
43. *Aquilegia virgata*
 in umbris alpinis Gavresii dicti i pra di Gale Cl. Dama
44. *Helleborus hyemalis*
 H. flor. folio incidente fl. luteo. / Aconitum unispolium bulbosum
 Copiosissime nascitur in vineis et collibus Monti Regali i pra di.
45. *Dianthus glaucus*
 in collibus Monti Regalensibus.
 D. fl. solitarii aut subolit. squam. calyc. lanceol. quaterni
 brevibus coroll. crenatis.
46. *Cucubalus italicus*
 c. calycibus clavatis, petalis semibipidis, panicula dichotoma
 erecta genitalibus declinatis caule erecto.
 Copiose ad Jesser circa Vico di Mondovi.
47. *Agrostemma Coronaria*
 A. tomentosa fol. ovato-lanceolatis petalis emarginatis coro-
 llatis serratis.
 Lychm. tomentosa Calyc. coccatis floribus longissime pedicatis
 loci siccis Monti Regali ubi circa loburent Montale & Frabosa.
48. *Viola Ruppii*
 V. caule erecto multiflora fol. ovato-lanceol. serratis
 in montibus Monti Regalensibus.
49. *Cistus pilosus*
 C. suffrutescens stipulatus erectiusculis fol. linear. subul. ciliatis
 incan. calyc. laevibus.
 passim inter Gavresio & Ormea.

The first of the month of January
was a very cold day, with a heavy
frost, and the wind was from the
north-east, bringing with it a
great deal of snow.

The second day was also very cold,
and the wind was from the north,
bringing with it a great deal of snow.

The third day was a very cold day,
and the wind was from the north,
bringing with it a great deal of snow.

The fourth day was a very cold day,
and the wind was from the north,
bringing with it a great deal of snow.

The fifth day was a very cold day,
and the wind was from the north,
bringing with it a great deal of snow.

The sixth day was a very cold day,
and the wind was from the north,
bringing with it a great deal of snow.

The seventh day was a very cold day,
and the wind was from the north,
bringing with it a great deal of snow.

The eighth day was a very cold day,
and the wind was from the north,
bringing with it a great deal of snow.

The ninth day was a very cold day,
and the wind was from the north,
bringing with it a great deal of snow.

The tenth day was a very cold day,
and the wind was from the north,
bringing with it a great deal of snow.

The eleventh day was a very cold day,
and the wind was from the north,
bringing with it a great deal of snow.

[Faint, illegible handwritten text, likely bleed-through from the reverse side of the page.]

[Marginal notes on the right edge of the page, including:]

- 30. ...
- 31. ...
- 32. ...
- 33. ...
- 34. ...
- 35. ...
- 36. ...
- 37. ...
- 38. ...
- 39. ...
- 40. ...
- 41. ...
- 42. ...
- 43. ...
- 44. ...
- 45. ...
- 46. ...
- 47. ...
- 48. ...
- 49. ...
- 50. ...

50. *Jedum Cappa*
J. foliis planis caule ramoso floribus paniculatis.
In ericetis umbratis in valle Tanari frequens.
51. — *Galioides*
Similis G. Cappa sed nobis distinguitur non procumbens ut prior
sed se erigit fol. inferior. 3. & 4. junct. deinde opposita prot.
bractea alterna vacua flor. etiam axillares sed proprios petal.
purpurascunt et arctata sunt teretifolia.
Supra Garresio loco dicto il Garbo della Luna.
52. *Ribes lva Crissa*
R. ram. aculeatis. Bacc. glabr. pedicellis bractea monophylla
in montibus Garresianis. Vidit Cl. Dana
53. *Pyrus Cydonia*
ad Septem in provincia Montivregali.
54. *Orchis anthropophora*
ophrys bulbis subrotundis Scapulo foliolo nectarii. Labio lineari
3 partito medio elongato bifido.
Inter Garresium & Ormeam.
55. *Fritillaria meleagris*
in pratis alpium Montivregalium.
56. *Alium europaeum*
Vulgare in umbris Montivregali.
57. *Buxus sempervirens*
Sylvatica facit inter Garresio & Ormea.
58. *Arum Dracunculae*
A. fol. pedat. folioli lanceolati integerr. superantibus spa-
tham pedice breviorum.
Circa Montaldo di Mondovi Synathium Cholinum.
59. *Agrostis miliacea*
A. petalo exteriori arista terminali recta stricta medioevi
Circa Garresio et Ormea
60. *Phromus inermis*
B. panicula erecta spiculis subaequalibus subulatis nudis submutatis
Circa Ormea lectus est.
61. *Staphisagria plicatulum*
locus flumen Costalicia Montibus Regaleis.
62. — *Spicant*
In montibus supra Roburent.

- Auctarii.
63. *Campanula latifolia*
Maxime frequent in pinguibus pratis di. Calotto.
 64. *Artemisia abrotanum*
prope Ceva & Garveto Mastrur.
 65. *Doronicum austriacum*
locis pinguibus & humidis; secus torrentes in montibus di. Calotto
& Frabosa.
 66. *3. beris nana*
3. herbacea fol. petiol. subrotund. dentatis flor. umbellatis.
In alpinis della Marchiera Monti Regali.
 67. *Cardamine hirsuta*
in sylvis ceduis subhumidis circa Casthus. di. Calotto paucim reperitur.
 68. *Arabis scabra*
Crescit frequent in alpinis di. Mondovi ubi la Marchiera versus Casthus.
 69. *Turritis stricta*
T. glabra lucens caule simplicissimo erecto silig. n. planis margine
attenuatis.
In pratis & paucis subhumidis; supra Calotto.
 70. *Sisymbrium barbarea*
S. fol. simpl. spatulato-ovatis; dentatis amplexicaulis nudis.
In agro Monvegalensi secus agros & locis siccis.
 71. *Papaver alpinum*
Vulgare in agris Monvegalensibus
 72. *Fumaria capreolata*
Inter la Jrea & Rastello & prope Ormea.
 73. *Busplenium petraeum*
B. cant. nudo fol. gramineis involucri & involucri distinct. polyphyllis.
In rupibus supra Ormea.
 74. *Chaerophyllum sylvestre*
abunde locis umbratis & pinguibus secus Sylva prope Casthus. Calotto.
 75. *Dianthus deltoideus*
D. flor. solitar. squam. calycin. lanceolat. binis corollis crenatis
repens.
In pratis montium Monti Regali.
 76. *Silene quadrifida*
locis rupestribus ad fontes in alpinis Monvegalensibus frequens.

[Faint, illegible handwritten text, likely bleed-through from the reverse side of the page.]

87. *[illegible]*
88. *[illegible]*
89. *[illegible]*
90. *[illegible]*
91. *[illegible]*
92. *[illegible]*
93. *[illegible]*
94. *[illegible]*
95. *[illegible]*
96. *[illegible]*
97. *[illegible]*
98. *[illegible]*
99. *[illegible]*
100. *[illegible]*

77. *Cucubalus alpestris*
C. dichotomus foli. linear. Calycibus petaliformibus inflatis. Petal.
 bispertitis.
 In alpibus d. Montis e. ruppium fissuris locis umbratis.
78. *Cirsium lunulatum*
 In montibus supra Ormea in jugis la Rarchera.
79. *Alchemilla nigra*
 Oritur et frequens est in alpebus Montis Regalis.
80. *Fritillaria involucreta*
 supra Ormea locis siccis & vespertinis max. spectantibus.
81. *Juncus filiformis*
 passim locis protuberantibus alpium Monregalensium.
82. *Phleum michelii*
 locis aridis et siccis supra Ormea.
83. *Festuca heterophylla*
 locis montanis siccis montium d. Montis.
84. *Avena festiva*
A. panicula patente locis densis fructibus festibus floribus
 duobus aristis.
 In agris avenae Montis Regalis.
85. — *bromoides*
A. subspicata spiculis binatis altera pedunculata arista divaricata ca-
 lycibus octofloris.
 In summis alpium Monregalensium soli expositis.
86. *Carex Bellardi*
 locis alpestribus solique patentibus alpium Monregalensium versus
 Carlin.
87. — *elongata*
 in pratis spongiosis alpium d. Carotto.
88. — *carecens*
C. spiculic. subrotundis remotis sessilibus obtusis androgynis cap-
 sulis ovatis obtusiusculis.
 in pratis spongiosis supra Carotto.
89. *Polypodium limbo-permum*
 in alpebus Monregalensibus.
90. *Saxifraga lingulata* Append: cit.
 frequens locis alpinis Monregalensibus.

91. *Boletus Frondosus*

St. sub caulis frondosus cespitosus sicut frondibus imbricatis planis
li. reflexis poris albis.

ad caudices arborum vehicularum & suspendam quandoque acquirit
magnitudinem a Montegalentibus Barberis dicitur.

Cunei die 5. Junii 1800.

Handwritten text at the top of the page, possibly a title or header.

Handwritten text block, likely the first paragraph of the letter.

Handwritten text block, likely the second paragraph of the letter.

Handwritten text block, likely the third paragraph of the letter.

Handwritten text block, likely the fourth paragraph of the letter.

Handwritten text block, likely the fifth paragraph of the letter.

Handwritten text block, likely the sixth paragraph of the letter.

Handwritten text block, likely the seventh paragraph of the letter.

Handwritten text block, likely the eighth paragraph of the letter.

Handwritten text block, likely the ninth paragraph of the letter.

Handwritten text block, likely the tenth paragraph of the letter.

Handwritten text at the bottom of the page, possibly a signature or footer.

1. Valuation
2. 1st. m.
3. 2nd. m.

4 fol.
3 in m

9 — 22
N. fol. in
Zach.

4. Maria b. Ziegler

Salvia
fol. ob
c. par. v
vid. l. m.
3. m. a.

6. — ~~by~~
~~by~~
~~by~~
pictures
3m. 01.

Geny C
Palacio
Miami
J. Nelson
Wato
in Agr

Mallotium
in larce

Stachyran
verhüllter
Frequenz in

— as
valicilla
function of

Plantae quae in Liguria Crescunt adnotante
C. Allionio in sua Flora pedemontana.

1. *Valeriana rubra.*
N. flor. monandr. caudat seu calcaratis foliis lanceolatis glaberrimis integerrimis.
In maceris ligurum.
2. *_____ angustifolia*
N. fol. integr. longissimis & angustissimis flore calcarato donato.
In maceris ligurum ubi vocatur Corinella
3. *_____ celtica*
N. fol. integerr. radical. ovatis caulinis linearibus obtusis.
In liguria alpina sterilibus non procul a Javone urbe.
4. *Rubia hirtorum*
In liguria. etiam circa Uneliarum.
5. *Salvia viridis*
S. fol. oblong. crenat. corollatum galea semiovalata calycibus pa-
chis reflexis bracteis cordatis acuminatis sensim decrescentibus
viridibus in spicam tetragonam digestis.
In arvis Uneliensibus
6. *_____ hispanica.*
~~*S. fol. ovatis crenatis dentatis verticillis subnatis stylo corollae la-*~~
~~*borio inferiore incumbente. S. fol. ovatis petiol. utriusque mucronatis*~~
~~*spicis imbricatis calycibus hispidis*~~
In Oliveis Uneliar.
7. *Sideritis Romana*
Genus Calycis subspidus galea erecta plerumque bipartita barba hispida
laciniis lateralibus acutis, media subrotunda stigma superius cili-
atum inferius membranaceo involutum.
S. herbacea decumbens ebracteata calycibus spinosis labio superiori
ovato.
in Agri Uneliar.
8. *Marrubium vulgare*
in Sarcotis liguria
9. *Stachys annua*
S. verticill. 6. flor. fol. ovatis lanceolatis. hirsuta. levig. petiol. caule erecto.
Frequens in liguria
10. *_____ avensis*
S. verticill. 6. flor. fol. obtusis nudiusculis. coroll. longitudine calycis. caule debili.
Uneliae secus arva.

11. *Myosotis apula*
M. coroll. bic. calyc. superant. semin. nud. latibus foliis lanceolatis hisp. ^{racem. pilosis}
In agro uneliensi licet sterilibus & semina.
12. *Covis monipulensis* in Montibus Ligurum.
13. *Antirrhinum bellidifolium*
A. fol. radical. lingulat. dentatis lineatis caulinis parvis integerrimis
in Liguria Collibus Circa Oneliam
14. *Scrophularia Vernalis*
pariam in Liguria locis sylvestribus & umbratis
15. *Plantago psyllium*
in aridis Sabulosis Liguria.
16. *Samolus Valerandi*
ad illigua Circa Oneliam.
17. *Lysimachia tenella*
L. fol. ovata acutiuscul. peduncul. foliis longioribus caule repente.
unelias secus torrentes & ad fontium scaturigines.
18. *Hyoscyamus albus*
H. foliis petiolatis sinuatis obtusis floribus sessilibus
In Liguria
19. *Hyoscyamus aureus*
H. fol. petiolatis dentatis acutis flor. peduncul. fructibus pendulis
ad olivos unelias.
20. *Verbascum phlomisoides*
V. fol. ovata utriusque tomentosis inferioribus petiolatis ligulate bicoll.
in aridissimis & siccioribus Liguria.
21. *Verbascum sinuatum*
V. fol. radical. pinnatifido-repandis tomentosis caulinis amplexicaulis
nudiusculis ramis primito oppositis.
ad Maris litora & Javona frequent.
22. *Convolvulus althaeoides*
C. fol. cordat. sinuat. sericeis foliis repandis pedunculis bifloris
in Liguria ad pagum Villatata.
23. *Convolvulus cheorum*
C. fol. lanceol. tomentosis flor. umbellatis calycibus hirsutis caule erecto.
Circa Oneliam.

24. *Convolvulus Soldanella*
C. fol. reniform. pedunculis unifloris
in maritimis Uneliae.

25. *Erica Cinerea*
E. antheris (vittatis) styloque inclusis corollis ovatis foliis ternis floribus
glomeratis
in montibus Uneliae.

26. ——— *Scorpiaria*
E. antheris bicornibus inclusis corollis campanulatis longioribus foliis
ternis patentibus ramis albis.
Circa Albenga.

27. ——— *arborescens*
E. antheris bicornibus inclusis corollis campanulatis longioribus foliis quaternis
patentissimis caule subarborescente tomentoso.
In collibus maritimis. Liguaria ut Circa Tagia.

28. *Cotyledon Umbilicus Veneris*. *Decandria Pentagynia*
Gen. Calycis 5-pidus Corollae campaniformis limbo 4-vel 5-fido squa-
mae nectariferae 5. ad basin geminis Capsulae 5-fol. carnosae succulentae.
C. fol. cucullato-peltatis serratis-dentatis caul. ramis. flor. erect.
in Maceris Liguarium frequent.
Caul. semiped. subnud. fol. altern. flor. terminali racemosa. Coroll.
albida subflavesc. aut purpurea.

29. *Olea Europaea*
in Liguaria maritimis.

30. *Momordica Elaeagnum*
M. pomis hispida cirrhis nullis
in Liguaria calidioris.

31. *Vitex agnus Castus*
Gen. *Didynamis angiosperma* Cal. 5-pidus corollae vixenti 2-labiate
lab. super. emarginato infer. 3-fido fructus est bacca 4-locularis loculis
monospermis.
In collibus maritimis prope Albenga.
V. fol. digit. serratis. typic. verticillatis
Arbor ramis folisque subtus incanis foliolis 5-7 lanceol. glabris.
Coroll. albo-purpurea vel tota alba.

32. *Globularia alyssum*
G. caule strigoso fol. lanceol. 3. dentatis, integrisque
in maritimis Uneliae.

33. ~~Glebularia~~ Xanthium spinosum
X. spinis ternatis foliis trilobis.
ad Charit. litus occurrit.
34. Echinops Sphaerocephalus.
E. capitul. globos. fol. sinuatis. pubescentibus.
in Collibus liguriae non infrequens.
35. Carduus linchus
Serratula fol. lyrato pinnatifidis pinnat terminali maxima flor-
culis confusis.
in Sylvis liguriae non infrequens est.
36. Cirsium medium X. p. Tenda.
in montibus circa Ugentiam.
37. Castanea lanata
C. caul. l. p. calyc. sanguin. terminal. primo aciculis sessil.
ad maris litus locis siccis.
38. Centaurea Galachites
C. calyc. setaceo-spinos. fol. decurrent. sinuatis. spinosis subter
fomentis.
In Collibus Ugentiae.
39. ——— elegant
C. caul. simpl. fol. indivis. linear. aciculis uniflor. & flor. terminal.
in Vineis Ugentiae.
Varietas praecedentis. Alti Auct.
40. Artemisia abrotanum
in pascuis & sterilibus locis collium Calidiorum liguriae.
41. ——— lobelii
in Collibus liguriae non infrequens.
42. Chrysocoma linolyvis X. p. Monrey.
in liguria saepe nascitur.
43. Conyza sordida
C. fol. linear. integerr. pedunc. long. 3 flor. caul. suffrutescente.
ad rupes & vetustos muros in Comitatu Ugentiensi.
44. Muphitalmum spinosum
M. calyc. acut. folior. fol. altern. lanceolat. amplexicaul. integer.
caul. herbac.
in liguria parce invenitur.
45. ——— salicifolium
M. calyc. ~~obtus. folior. pedunculat.~~ fol. altern. lanceolat. suberr. villis.
herbac. Abunde in liguria.

o/o.

integral

6/err. V
General

no. 1164 in *Xanthium Michxolium*

[illegible]

46. *Achillea ligustica*
in collinis supra oppidum et Cairo.
47. *Aster acris*
A. fol. lanceol. linear. strict. integerr. panic. floribus corymbois
subligatis pedunculis foliolis.
in pratis unelicis.
48. — *bipennis*
Bruta fol. oblong. decurrentibus denticulatis floribus corymbis ter-
minalibus subsessilibus.
in Liguria.
49. *Erigeron graveolens*
E. ram. lateral. multiflor. fol. sublinear. integerr.
in collibus maritimis unelicis.
50. *Senecio Sylvaticus*
S. coroll. revolut. fol. pinnatif. denticulat. calic. corymb. erect.
in Liguria.
51. *Cineraria maritima*
C. flor. panicul. fol. pinnatif. tomentis. incan. sinuatis. caul. stratis
ad Rhodum unelicis.
52. *Catananche carulea*
Gen. Cal. imbricatus. Receptac. paleaceum pappus aristatus
calyculo 3-seto.
C. squamis calyc. inferior. ovatis.
abunde in sterilibus Liguriis.
Caul. 2. cubit. teret. subimpt. fol. altern. sessil. linear-lanceol.
subdent. 3-nerv. pedunc. termin. 1. flor. petal. coroll. erect.
villos.
53. *Picris hibernica*
Leontodon. fol. pinnatif. stratis. Calyc. acut. hirsut.
in pratis Liguriis.
54. *Hieracium tomentosum Aurigala lanata L.*
in Liguria.
55. *Crepis pulchra*
C. fol. sagitt. denticul. caul. panicul. flor. pyramid. glabris.
in Vineis & Uliveti unelicis passim.
56. *Sonchus tenuissimus*
abunde in Liguria.
57. *Scorzonera graminifolia*
S. fol. linear. ensiform. integris cavinatis
secus agros unelicis.

58. *Tragopogon picroides*
T. calycibus monophyllis; corollis brevioribus aculeat. fol. runcinat.
denticulatis.
in Olivetis Unelia.
59. *Epimedium alpinum* V. plant. Monregal
in Liguria.
60. *Alyrium maritimum.*
Clypeola perennis silicul. biflorulav. ovat. disperm. fol. linear.
in arenosis maritimis Unelia.
61. *Linaria halmifolia* V. pl. Valder.
in collibus Unelia editionibus non tamen ita quicquid sunt proceri.
62. *Sinapis maritima* V. pl. Vinad.
Nascentur ad ora maritima Unelia.
63. *Hyperic. lychnitis*
H. caul. subhispido erecto valvis latius. Aliguae.
in Valle Unelia.
64. *Ruta graveolens*
in principatu Unelia. Varietas minor frequentior est.
65. *Euphorbia spinosa*
E. umbella subsida simplici involuelli ovali primariis bi-
phyllis fol. oblong. integerr. caul. pubescent.
ad Ruper in Liguria nascitur.
66. *_____ epythymoides*
E. umbella bifida, bifida involuelli. ovata fol. lanceolata. obtus. subhy-
villosa.
Locis incultis & ad ora sylvarum Liguria non infrequens.
67. *_____ dendroides.*
E. umbella multipida dichotoma involuelli subcordati primariis
triphyllis caule arbores
ad scopulos mari proximos prope Uneliam & Alarciam.
68. *Papaver hybridum*
sp. caputis subglobosis tomentis hispidis caul. folior. multiflor.
inter segetes. Unelia.
69. *Trifolium ochroleucum.*
T. caul. hirsut. scabr. fol. integerr. moll. spic. subvillos. ochroleuc.
Locis apricis collinis, Liguria, frequent est.
70. *Lobus edulis*
L. leguminibus subrotundis gibbis incurvis
in pascuis sterilibus Unelia.

2/1. *Colus ornithopodoides*.

1. Colus ornithopodoides - *1. Colus ornithopodoides*

2/1.

1. Colus ornithopodoides

58. *Tragopogon microides*

T. calycibus monophyllis corollis brevioribus aculeatis. fol. runcinatis.

11. 10/11

59

60

61

62

63

64

65

66

67

68

69

70

0/1. *Lobelia ornithopodoides*.

Lobelia ornithopodoides L. f. 115 2. 0

h/.

in Liguria humiliter? planta.

58. *Tragopogon picroides*
T. ciliatus monstrosus, corollis brevioribus aculeis. fol. runcinat.

59

60

61

62

63

64

65

66

67

68

69

70

71. *Lobelia*
L. longifolia
 in l.

72. *Dory*
 in l.

73. *Medicago*
M. pauciflora
 3h 1/2

74. *Ononis*
 in l.

75. *Cytisus*
 par

76. —
 in l.

77. *Hedysarum*
H. foliosum
 3h 1/2

78. *Onobrychis*
Hedysarum
alysoides
 in l.

79. *Vicia*
V. longifolia
 in l.

80. —
 in l.

81. *Ornithoglossum*
O. foliosum
 ad ag.

82. *Sparganium*

83. —
L. foliosum
 in l.

84. *Cercis*

85. *Athyrium*

71. *Oxalis corniculoides*.
l. legum. subternat. arcuat. compress. caulis diffus.
in Liguria.
72. *Dorychnium Monspeliense*
in Liguria abunde.
73. *Medicago marina*
M. pedunc. racemos. legum. Coch. spinos. cauli. prostrat. b. toment.
In sterilibus litoreis maris proximis.
74. *Ononis Columna*.
in Liguria frequenter obviam venit.
75. *Cytisus decussatus*
passim in Liguria.
76. ——— *argenteus*
in collibus Liguriae.
77. *Hedysarum spinosissimum*
H. fol. pinnat. legum. articulat. aculeat. toment. cauli. diffus.
In agro unelienti locis sabulosis mari proximis & secus arva.
78. *Onobrychis sarcocolla*.
Hedys. fol. pinnat. legum. l. spinos. sulcat. toment. corollae
altissimis scapis subradicatis.
in sarcocollae apricis uneliae.
79. *Vicia Seppium*
V. legum. pedicell. subquatern. erect. foliol. ovata. integerr. escheno-
vibus decrescentibus
in editionibus Liguriae montibus A. Jura.
80. ——— *harbonensis*
in dumetis Liguriae.
81. *Oxalis corniculoides*
O. fol. ternatis subternatis impari maximo.
ad agros macilentos in Liguria.
82. *Spartium juncum*
in Liguria.
83. ——— *spinosum*
S. fol. ternatis v. impari angulatis spinosis
in Olivetis uneliae.
84. *Cercis siliquastrum*
in Liguria maritima
85. *Astragalus mollis*
in Liguria humilis? planta.

96. *Eryngium maritimum*
l. fol. radical. subrotund. plicat. spinos. capitul. pedunculat.
paleis micropidatis
in sabulo ad maris litus.

97. *Platinaca oppositifolia*
p. fol. bipinnatis foliol. oblique cordatis petalis paleaceo (trigori).
Uetia locis umbrati & montanis ruderatibus.

98. *Ferula nodiflora*
Gen. Semen magnum oblongo-ovatum 5. liliatum Cinchumala magna
Solida non membranacea.
F. foliol. appendiculatis umbellis sessilibus
ad Vineas in collibus Uetiae.

99. *Pencedanum officinale*
p. foliol. hier. bipinnatis filiformibus linearibus
in montibus liguria.

90. *Anerthum graveolens* fructibus compressis
ad maris litus prope Uetiam.

91. *Angelica Levisticum*
Ligusticum fol. multiplic. foliol. superm. incis. flor. luteolis
inter Savonum et Genuum spontaneam habent liquores ejusque caulibus
& folia edunt inter Apii Graveolentis.

92. *Coriandrum tereticulatum* fructibus didymis
passim in agris Uetiae

93. *Oenanthe spinipinelloides*
Gen. fl. radiati disco sterili petala inaequalia fructus ovatus calyce
inligui 5. dentato persistente coronatus sem. ovat. ^{non} acuminat.
thialium inter plantarum
O. foliol. radicalibus cuneatis filis caulibus integr. linear. longioribus
canaliculatis.
In liguria praesens occurrit.

94. *Bupleurum rotundifolium*.
p. involucell. univers. nullis fol. persiliat.
In agris collium calidiorum liguriae passim speciatim Uetiae.

95. ——— junceum
B. caule erecto paniculato fol. linear. involuc. 3phyll. involucell. 5phyll.
Jecur agris in liguria.

96. *Faniculum officinale*
in Provincia Uetiae.

97. *Bunium minus*.
p. fol. unifolmiat involucro polyphylo fructibus subcylindricis apice cili-
sistibus stylis reflexis deciduis
In edioribus montibus liguriae.

98. *Crithmum*
 Gen. Flor.
 Coharent
 C. folia
 ad Auph

 99. *Caucalis*
 Scandisc
 in Vineis

 100. *Ronunculus*
 R. semini
 in arvis

 101. *Corrigiola*
 Gen. Cal. 5p
 Lat. are
 Cant. prot
 Nipul. bir
 Congesti in

 102. *Potentilla*

 103. *Helicorhiza*
 in

 104. *Silene* Jovic
 l. fol. acut
 ad Mav

 105. *Oxy* cornic
 in mo

 106. *Reseda* l
 in

 107. *Tribulus*
 Cal. profor
 pitulum
 T. fol. sub
 in liguri
 Cant. pedat
 nata foliolis
 lavy fascicu

 108. *Quonimus*
 e. fol. acut
 in Edibov

 109. *Cichus* albid
 C. arboresc
 ferilibus
 in incultis

98. *Cisthmm maritimum*
 Gen. Flores aequales fertiles. Semen ovale intus planum tota superficie
 cohaerens dorso tribus obtusis costis percussio.
 C. fol. lanceolat. carnosis.
 ad Ruper mari proximis in muris. Vetusis Unelia.
99. *Caucalis nodosa*
 Scandisc seminibus subcylindricis hispida, caule hispido genicul. humidi
 in vineis & sterilibus figuris minime varia.
100. *Ranunculus parviflorus*
 R. seminibus mucatis fol. simpl. laciniat. acut. hirsut. Caule diffuso
 in arvis subhumidis Unelia.
101. *Corrigiola litoralis*
 Gen. Cal. 5 phyll. Stamina 5. petal. 5. Sem. 1. Wignetr. nud. calyc. content.
 Locis arenosis mari proximis.
 Caul. prostr. virens. Varnosi fol. altera. sessil. lanceol. integer.
 bipul. binat. acuta argentea. flores racemos. in ultimis varis
 congesti interd. arcuati petal. Oval. alba.
102. *Potentilla recta*
 in Liguria valde frequens.
103. ——— rubens
 in Calidioribus collibus Liguria.
104. *Helleborus hyemalis* V. pl. Monveg.
 in vineis et collibus Liguria.
105. *Silene Sericea*
 S. fol. acuminat. Succulent. hirsut. lucid. flor. carnea
 ad Mavis lites in Sabulosis inter Oneglia & Porto Maurizio.
106. *Oxy corniculata*
 in montibus Liguria ad iurigna frequens.
107. *Aerida luteola*
 in Liguria ad vias muros & rudrata.
108. *Tribulus terrestris* Decand. monogyn.
 Cal. profunde 5-fidus petal. 5. aequal. stili nullus. Stygma la-
 pitulum capsulae plures congestae aculeatae.
 T. fol. sub 6-jugis subaequalibus seminibus quadrangulis
 in Liguria
 Caul. pedat. & ultra prostrumb. ramos. villosus rubens. fol. abrupte pin-
 nata foliolis oblique lanceolatis ciliatis subus villor. flos et arcu-
 lare fasciculati peduncul. flavi
109. *Quonimus latifolius*
 e. fol. acute & argute serratis capsul. angul. acut. membranaceis
 in edibilibus montibus Liguria.
110. *Cistus albidus*
 C. arborescens erectissimus fol. ovato lanceolatis tomentosis incanis
 sessilibus subminuor
 in incultis & montibus collibus Unelia.

111. *Cistus guttatus*.
C. herbaceus erectissimus fol. oppos. lanceolatis brevibus. racem ebract.
C. flor. pallid. macula punicea in rugis
in pascuis dicitur sterilibus Liguria.
112. — *Pilosus*
in lignum montibus.
113. — *Vosens*
Urtica natrix.
a C. helianth. differt. caul. magis ramos. & fruticosa. fol. firmior.
subtus subincanis corollis roseis.
114. *Linum hirtutum*
L. calyc. hirsutis acuminatis dent. altern. fol. alternis, varicis oppositis.
non rarum in Liguria.
115. *Jedum Cespica*
J. fol. planis caule ramoso flor. paniculatis.
in collibus Liguria.
116. — *acre*
in dicitur pascuis Liguria.
117. *Rhamnus infectoria.*
R. spinis terminalibus flor. 4-fidis dioiciis caulibus procumbent.
in Liguria lectus est.
118. — *Lycioides*
R. spinis terminalibus foliis linearibus
in Liguria habuit Cl. Alton.
119. *Rhus cotinus*
frequentia in montibus Liguria.
120. *Prunus domestica*
pr. pedunculis solitariis fol. lanceolat-ovatis convolut. serrat. ram. multis.
in sylvis Liguria.
121. *Myrtus Communis*
Glen. calyce superius 5. vel 4. fidis petalis. vel 4. ovatis staminibus
plura calyce inserta stigma obtusum bacca olivaceo. coronata
umbilicata 2 vel 3. locularis
Myrtus minor Vulgaris
Savona reperitur.
Planta pubesca fol. busco acutiora opposita subsessilia ovata-acu-
minata laevia satore viridia odore petala alba.
122. *Azota rubiginosa*
A. germ. globosis petiol. aculeis. aculeis recurvis fol. subt. rubiginosis.
in collibus Liguria ad Viar, & sepe occurrit.
123. *Pyrus Cydonia*
ad saepe in Liguria.

124. *Orchis*
 125. —
 126. —
 127. —
 128. —
 129. —
 130. —
 31. *Epipactis*
 32. *Pancreaticum*
 33. *Allium*
 34. *Asphodel*

124. *Orchis*
 125. —
 126. —
 127. —
 128. —
 129. —
 130. —
 31. *Epipactis*
 32. *Pancreaticum*
 33. *Allium*
 34. *Asphodel*

124. *Orchis Coriophora*
O. bulb. indivis. nectar. lab. 3. fid. reflexe. crenat. corum brevi.
petal. connivent.
in prati/ siccioribus collium Liguriae non infrequens.
125. — *hircina*
O. rad. subrotund. label. longiss. 3. spathato plicato.
Sylvos. et apricis locis collium Liguriae. sed non frequens.
126. — *muticaria*
O. rad. subrot. label. sericeo bitulro brachiol. brev. subulat.
in sterilibus graminosis collium Liguriae.
127. — *avachnites.*
O. rad. subrotund. label. holosericeo emargin. appendiculato.
abunde in collibus Liguriae quandoque cum praeced. (venit).
128. — *lingua*
sevapina bulbis subrotundis nectar. lab. 3. fido acuminato petalis
longiore glabro.
in prati & pratis subhumidis Liguriae frequens.
129. — *anthropophora* V. pl. Monvegat.
in dumetis umbrosis unctis quandoque occurrat frequenter ad
oppidum monte d'Alto.
130. — *fusca*
O. rad. subrotund. spica long. label. 4. fido brachiol. angust. ciliolatis labis
sevapinis.
In collibus Ligurum nascitur.
131. *Epipactis aethiva.*
O. bulbis fasciculatis filiform. nectar. lab. ovat. integerrimo caul. aphyll.
in dumetis et sterilibus Calidivum collium Liguriae.
132. *Pancreaticum maritimum.*
Gen. Calyc. est spatha Nectarium infundibuliforme 12. fid.
fructu impositum stamina sustinent Stamina 6. & 6. fid. petala.
sp. spatha multiflora petalis planis filis lingulatis.
ad Vesper prope litus maris ubique frequens
Fl. magn. petal. lanceol. alba sub hyl. linea viridi.
133. *Althium roseum*
A. caul. planifol. umbellifero umbella fastiginta, petalis emar-
ginatis stamin. breviss. simplicibus.
In Olychis et arvis unctis.
134. *Asphodelus fistulosus*
A. caul. nudo. fol. strictis subulatis striatis subfistulosis.
in Liguriae montibus in sona proximis

135. *Scilla maritima*.
S. nudiflora bracteis reflexis
 ad maris litem arenosa & suppet inhabitat.
136. *Asparagus acutifolius*
A. caule inermi angulato fruticoso fol. acinis vix rigidis perennantibus mucronatis aequalibus
 in principatu Unelias.
137. *Lythrum hyssopifolia*.
L. fol. alternis linearibus fl. descendis
 in humidis & salubris Liguria.
138. *Adonis flammea*.
A. fl. octopetalis fructibus subcylindricis calyc. hirsutis
 in collibus calidioribus Liguria.
 Floretestate petala flammea macula fere nulla.
139. *Pinus sylvestris*
 in montibus montanis & collibus Liguria.
140. *Juniperus oxycedrus*
J. fol. ternis patentibus mucronatis bacca brevioribus.
 in sylvis appicis versus Uneliam.
141. *Phanicea*
J. foliis ternis oblitteratis imbricatis obtusis.
 frequent in collibus Unelias.
142. *Alviperia vicia*.
A. caule herbaceo fol. incan. serratis fruct. 4. angul. dentato vix compresso.
 In Liguria frequent est.
143. *Hedysarum hirsutum*
 in Liguria frequent.
144. *Salsola Kali*
*Calyc. profunde 5-fidus quodammodo coloratus pedicellus stylus lim-
 plere semibifidus Stamina 5. Capsula globosa serena. 1. cochlearina*
J. decumbens herbacea fol. subulat. spinol. scabris calycibus
marginalis arcuatis.
 in maritimis prope Uneliam.
 Planta spinosa fol. crassa calyc. cinch. margine foliaceo stylus 3fid.
145. *Juncus acutus*
J. culmo subundo teret. mucronato terminali involucri 2-phyllo
 spinoso.
 in locis maritimis Unelias.
146. *Buncus sempervirens*
 in montibus Liguria ad suppet.

[Faint, mostly illegible handwritten text in a cursive script, likely a historical manuscript or letter.]

in figura ad littera man.

26

26

lat. fin.
h. l. i. h. n.
c. 6. 4.

Stylus

2. p. 4. 4.

[Faint, mostly illegible handwritten text in a cursive script, likely a botanical manuscript. The text is arranged in several horizontal lines across the page.]

in molibus figura ad v. p. 1.

117. *Artemisia*
A. fol.
in lob.
 118. — *h.*
A. fol.
in lob.
 119. *Viscum*
Gen. Ch.
calyci a
globosa
V. fol.
Unelia
Fol. op.
abou-va
 120. *Astrum a*
A. a. cov.
in lob.
 121. *Nardus ar.*
U. calyc.
ad
 122. *Alopecurus*
Ch. culmo
in pua
 123. *Rottbolla*
Gen. locusta
sed glum.
lig. tum.
A. spica bere
Unelia
 124. *Lolium ten.*
L. spica mus.
in pascuis
 125. *Aira canere*
A. fol. seta
in debul.
 126. *Poa rigida*
P. panicula
abunde in
 127. *Bromus ligu.*
Gramen Brom.
in figura a

147. *Ruscus aculeatus*
R. fol. supra florifer. nudis
in collibus Liguriae.
148. — *hypoglossum*
R. fol. supra florifer. sub foliato.
in lacunis in cultis Liguriae.
149. *Vitium album*. Dioecia tetrandria.
Gen. char. Cal. 4. part. segment. ovat. Coroll. 0. filam. 0. antherae
calyci adnatae. Fam. Cal. 4. phyll. superius Coroll. 0. Styl. 0. Bacca
globosa glutine plena 1. sperma semina cordata compressa.
V. fol. lanceolatis obtusis caule dichotomo spicis axillaribus.
Umelia et locis maritimis frequens et parasitica planta
Fol. opposita sessilia integerrima crassa dura sub lente. Baccae
albo-rubrae nitidae.
150. *Atrum aviculatum*.
A. acaule fol. cordat. oblongis spathae apertura ovata.
in collibus Umelia.
151. *Nardus aristata*
N. calyc. aristati.
ad Vini in Liguria.
152. *Alopecurus bulbosus*
A. culmo erecto spica cylindrica radice bulbosa
in pratis ludit Liguriae.
153. *Rottbolla incurvata*.
Gen. locustae uniflorae flexuosi caulis scrobibus appressis. Calyce 2. glumis
sed glumis iuncta se possit ita ut 1. glumis calyce videatur. folliculis
biglumis.
R. spica tereti subulata gluma calycina subulata appressa 2. partita.
Umelia in agris muni proximis.
154. *Lolium tenue*.
L. spica multica tereti spiculis trifloris
in pascuis Liguriae.
155. *Aira canescens*.
A. fol. setaceis summo spathaceo panicul. inferne obsovente.
in sabulis sterilibus Liguriae.
156. *Poa rigida*
P. panicula lanceolata subramosa secunda ramul. altern. secundis.
abunde in Liguria.
157. *Bromus Ligusticus*
Gramen bromoideae festucae tenuique panicula ardens
in Liguria ad litora maris.

158. *Elymus arenarius*
L. Spica erecta arcta calyc. tomentor. florculo longioribus
in arena ad maritima.

159. *Agropyron ovata*
L. Spica ovata calyc. omn. brevioribus
locis siccis in Liguria ubique

160. *Andropogon hirtum*
L. Panícula spica conjugatis calyc. hirsutis.
Frequens in lina locis aridis & siccis.

161. *Scirpus mucronatus*
L. Culmo triangulo nudo acuminat. spic. conglomer. sessil. lateralibus.
in maritimis lina.

162. *Schoenus mucronatus*
L. Culm. teret. nud. spicul. ovat. fasciculat. involucri sub 6. phyllo fol.
canaliculatis
In maritimis lina.

163. *Adiantum Capillus Veneris*
L. fol. decompositis foliol. altern. pinn. ~~diversis~~ cuneiformibus
lobatis ped. cellatis.
Vulgatum in pratis & rupibus Liguria Colliumq. maritimorum.

164. *Fucus serratus*
Chalc. Vesiculae villis intertextae Form. Vesiculae ad-
spere granis immixtis apice prominulis semina soli-
taria substantia coriacea Vesiculosis.
F. fronde plana dichotoma serrato dentata fructificatio-
nibus terminalibus tuberculosis
In mari lina.

165. ——— *Vesiculosus*
F. fronde plana dichotoma integerrima Vesiculis
axillaribus geminis terminalibus tuberculatis
in mari lina.

166. ——— *Nodorus*
F. fronde compressa dichotoma fol. dich. integerr.
Vesiculis annatis solitariis dilatatis
ad mare lina.

167. ——— *Selaginoides*
F. frond. filiform. Vam. dichot. subulatis
alternis basi vesiculosis
ad lina lina.

168. *Fucus* ...
 169. ...
 170. ...
 171. ...
 172. ...
 173. ...
 174. ...

Fucus

175. ...
 176. ...
 177. ...
 178. ...
 179. ...
 180. ...

68. *Fucus*
 169. —
 170. —
 171. —
 172. —
 173. —
 174. —
 175. —
 176. *Ulva*
 177. —
 178. —
 179. —
 180. —

168. *Fucus fasciculaceus*

f. fronde filiformi ramorissima vericulis ovalis terminatis
foliolis multipartitis obtusis apice fructificantibus
in Rupibus litoris Uneliae.

169. — *abrotanoides*.

f. fronde ramosa capillari multipida fasciculata vericulis glo-
boris.
in mari Uneliae.

170. — *acinarius*.

f. caule filiformi ramoso fol. linear. integerr. fructificat. globosis
pedunculatis
in mari Uneliae.

171. — *volundus*.

f. teres subdichotomus uniseriis ramis obtusis verrucosis
in mari Uneliae.

172. — *muscoideus*.

f. fronde tereti ramorissima ramis sparsis spinis mollibus alternis.
in mari Uneliae.

173. — *longissimus*.

f. teres rubens minutus ramosus in longum protensus
in mari Uneliae.

174. — *confervoides*.

f. subdichotomus teres filiformis ramoriss. ram. inaeq. setaceis.
in mari Uneliae.

175. — *plumosus*.

f. caule filiformi subcompresso ramoso frondibus lanceolatis
bipinnatis plumosis (purpureo) coloratis
in mari Uneliae.

176. *Uva intestinalis*.

Gen. Fructific. in subrotund. membranacea longa tubulosa
vel Campanulata larca diaphana.
U. tubulosa limplex intestinorum figura
in mari Uneliae.

177. — *compressa*.

U. tubulosa ramosa compressa
in mari Uneliae.

178. — *lactuca*.

U. palmata prolifera membranacea ramentis infern. angustatis
Fucus fronde imbricata frond. sparsialibus erectiusculis sinua-
to laciniatis obtusis Lactucae similis planta.
in mari Uneliae.

179. — *linga*.

U. fronde oblonga bullata
in mari Uneliae.

180. *Lycopodium* tuber.
in collibus liguriae.

181. *Aristolochia Clematidis*
in arvis subhumidis, liguriae.

182. *Statice luteola*
St. caepodichotomo ancipiti fol. ovat. mucronat. floribus
aggregatis.
Limonium fol. lanceolat. mucronat. radicalibus caul. ancipiti
dichotomo stipulis simplicibus.
Occurrit circa Ventimiglia

183. *Artemisia camphorata*
A. fol. petiol. palmato-multifidi, linearibus superveni simplicibus
calycibus angulosis receptaculo hirsuto.
Nascitur frequenter in Liguria locis superstitibus.

Cunei die 6. Junii 1800.

Hodie ab Exercitu Coelaveo in Cunei celebrata est deditio
Genuae subit principis Liguriae.











[Faint, illegible handwriting covering the upper two-thirds of the page. The text appears to be a continuous paragraph or a list of items.]

[Faint, illegible handwriting in the lower third of the page, separated from the upper section by a horizontal line.]

1. Velle
/ Ma
C. P. A.

2. —
V. ma
in u
C. A. P.

3. —
in u

4. —
V. /
in u
in
V. ma
V. ma
C. A. P.

5. —
C. A.
V. ma
V. ma
B.

6. Galium
G. fol.
Nico

(*) Mo.
in u
nat. facit
et alibi
p. centur
familiae
vult, et
Cuneis
parare
M. P. A.

Plantæ Nicæenses, & Comitatus quarum locus in Flora pæd. natalis designatur. (*)

1. *Valeriana rubra*. V. pl. ligur.
frequens ad muros. in vineis agri Nicæensis.
Circa Jaspello ubi appellatur *Cornacina*.
2. ——— *Calcitrapa*.
V. monostemon fol. im. integr. superioribus pinnatis
in umbrosis agri Nicæensis.
caules cubitales ramosi folia humifusa corolla dilute rubra.
3. ——— *tuberosa*. V. pp. Monnegal.
in Comitatu Nicæensi specialim supra *Cornacina*.
4. ——— *Jascanilis*.
V. fluv. triandris fol. subdentatis radicalibus ovatis caulinis linearilanceolatis.
in montibus Nicæensibus.
Planta nuda caul. spitham. simpl. fol. subtrinatis. ad ora obsolete pilosa radical. 4. caul. 2. opposita v. n. brachiatis p.
dunc. opposit. fl. panicul. corol. alba.
5. ——— *Coronata*.
V. caul. dichotomo fol. pinnatifidis sem. Corona radiatum dentata
dentibus 6.
3 h. avui Nicæensibus.
6. *Galium palustre*.
G. fol. quatern. obovat. inæq. caul. diffus.
Nicæe semper pascuis & humida loca.

(*) *Notandum*. Non omnes stirpes quas Celeber. Aldione in sua Flora pædemontana Comitatus Nicæensis indigenas facit hic decententur. Nimirum enim liber increveret, et alibi descriptæ plantæ denuo enumerarentur. Ideoque relincentur Stirpes eæ, quæ in pædemontio frequentes, et familiares sunt, vel in hostis Culta omnibus innotuerunt, & eæ quoque, quas ego vel in agro Patrio, vel Cuneensi jam observavi, quasque mihi velim comparare necessum non est, ut ad Comitatum Nicæensem propicietur.

7. *Galium uliginosum*
G. fol. sem. lanceolatis retrorsum serratis aculeatis mucronatis rigidis
 vallis fructu majoribus.
 in agro Nicaensi ad viv. loc.
8. — *Jascale.*
G. fol. sem. obovatis obtusis caul. ramosis. procumbenti-
 in ed. litoribus montibus S. Stephani Nicaensis.
9. — *purpureum*
G. fol. verticill. linear. setaceis pedunc. capillar. fol. longior.
 abunde in agro Nicaensi.
10. — *lucidum*
G. fol. 6. rigidis obscure-vir. flor. e summo caul. prodeuntibus
 ubique in agro Nicaensi. locis siccis.
11. — *cinerum*
G. caul. subligrolo fol. 6. long. elliptic. rigid. serrat. aculeat. var-
chibus ovatis laevibus albescentibus
 in vineis Savorgii & Sospitellii.
12. — *tenuifolium.*
G. fol. 6. subrigid. diffuse ramos. flor. alb. e summo caule prodeunt.
 in collium Nicaensium duneis.
13. — *murale*
G. fol. 6. linear. ram. simplicibus fructibus hispido-suberentibus.
therardia fol. florat. binis oppositis binis floribus.
 ad aridas rupes litoris Nicaensis.
14. — *maritimum*
G. fol. octon. lanceol. - linear. peduncul. 1. flor. caul. vespert. luto.
 In agro Nicaensi collectum est.
15. — *saccharatum*
G. sem. escarpent. florul. lateral. marcul.
 Circa Nicaeam provenit.
 Vera Varietas Valantia affavine L.
16. *Valantia muralis.* Coroll. 4. fida sem. tridentata.
fl. flor. marcul. tripartitis hermaphroditi germine glabro
insidentibus.
 in siccis & arenosis litoris Nicaensis.
 Planta glabra caul. procumb. fol. verticill. 4. Corol. e viridi-
 pallens fruct. asperi.
17. *Rubia tinctorum*
 ad muros & rupes ubique frequens in agro Nicaensi.

1. ...
2. ...
3. ...
4. ...
5. ...
6. ...
7. ...
8. ...
9. ...
10. ...
11. ...
12. ...
13. ...
14. ...
15. ...
16. ...
17. ...
18. ...
19. ...
20. ...
21. ...
22. ...
23. ...
24. ...
25. ...
26. ...
27. ...
28. ...
29. ...
30. ...
31. ...
32. ...
33. ...
34. ...
35. ...
36. ...
37. ...
38. ...
39. ...
40. ...
41. ...
42. ...
43. ...
44. ...
45. ...
46. ...
47. ...
48. ...
49. ...
50. ...
51. ...
52. ...
53. ...
54. ...
55. ...
56. ...
57. ...
58. ...
59. ...
60. ...
61. ...
62. ...
63. ...
64. ...
65. ...
66. ...
67. ...
68. ...
69. ...
70. ...
71. ...
72. ...
73. ...
74. ...
75. ...
76. ...
77. ...
78. ...
79. ...
80. ...
81. ...
82. ...
83. ...
84. ...
85. ...
86. ...
87. ...
88. ...
89. ...
90. ...
91. ...
92. ...
93. ...
94. ...
95. ...
96. ...
97. ...
98. ...
99. ...
100. ...

18. *Crucianella angustifolia*.
 Gen. Corol. 1. petal. intundibulata. tub. filiformi limbo
 unguiculato 4. fido calyce. 2. styli sem. 2. tenuia ob-
 longa flor. in spicis imbricatis folia verticillata.
C. erecta fol. dens linearibus flor. spicatis.
 Secus arva Collium Nicaeensis.
19. ——— *monspeliaca*
C. procumb. fol. acut. caul. 4. ovat. variet. linear. fl. spicat.
 Secus arva Coll. Nicaeent.
20. ——— *latifolia*
C. procumbens fol. 4. lanceolat. fl. spicat.
 in Comitatu Nicaensi locis arenosis non infrequens.
21. *Salvia officinalis*
 in Coll. incultis agri Nicaensis
22. ——— *horminum*
S. fol. obtus. crenat. bract. sum. steril. majoribus coloratis.
 Nicaentem agrum inhabitat Secus arva nascitur.
23. ——— *verbenaca*
S. fol. serrat. sinuat. levissim. Corol. Calyc. angustior.
 in Comitatu Nicaensi non infrequens est.
24. ——— *aethiopis*
S. fol. oblong. eros. lanat. Verticil. lanat. bractes recurvatis
sub spinosis.
 Circa Sorsello in Comitatu Nicaensi.
25. *Thymus vulgaris*
 in apricis saccis Collium Nicaeensis sponte nascitur.
26. *Salvia juliana*
S. Verticill. fastigiatis fol. linear. lanceolat.
 Circa Nicaeam in aridis
27. ——— *thymbra*
S. flor. Verticillat. hispida. fol. oblong. acut.
 in saccis Collium Nicaeensis
28. ——— *gracea*
S. pedunc. sub 3. flor. lateral. involucl. calyc. brevior.
 in agro Nicaensi.
29. *Thymbra verticillata* Vide St. Monveg.
S. fl. verticillatis.
 in agro Nicaensi.

30. *Lavandula stoechas*.
L. fol. lanceol. lineari. integerr. spicis comosis
in locis apertis Maritimi agr. Nicæensi.

31. *Sideritis Romana* Vid. pl. figur.
in arid. collibus phœacæ.

32. ——— *incana*
S. fruticosa tomentosa fol. lanceolat. lineari. integerr. flor.
bracteisq. dentat.
inter L. Skavene & Rocca talliada

33. ——— *hirsuta*
S. fol. lanceol. obtus. dent. pilos. bracteis dentat. spinos. caul.
hirs. decumbentibus.
in Comitatu Nicæensi ad vappes & colles.

34. *Stachys brevifolia*
S. hirsuta fol. obscure virentibus elliptico-lanceolatis, verticil.
10. flor.
in collibus Comitatu Nicæensi locis minime cultis.

35. ——— *hirta*
S. verticill. 6. flor. caul. prostratis corol. lab. superiori 2. pido
divaricato reflexo.
Habitat in agro Nicæensi.

36. ——— *arvensis* V. p. figur.
Nicææ secus arva.

37. *Phlomis lychmich*
Gen. Calyc. angulatus galea incumbens compressa hirta.
Ph. fol. lanceol. tomentol. floral. ovat. involucri. setae. lanat.
in Agro Nicæensi
Caul. cubital. Corol. vix calyce major lutea.

38. *Molucella frutescens*
Gen. Calyc. ampliat. campanulatus corol. latior spinosus.
M. calyc. in fundibuliformibus 5. pido corol. calyc. longioribus.
Fruticul. palmat. var. fol. floriferis dichotomis fol. mollis.
ovat. breviter petiol. utrinque 2-3. profunde serrata pilis variis
pubescentia nervosa venosa spin. areolaribus utrinque. solit. ternat.
ut in Berberide sed tant. ad florum verticillos. Cal. spinos. lo-
bratus corol. alb. galem lanatam.
Abunde inter Saorgio et Brelio et supra Ispello & vespum
filivis erit.

39. *Melissa cretica*
M. racem. terminal. peduncul. solitar. breviss.
Circum Lucerna.

longistylus
fol. pallid.
pil. var. /
solit. f. tenuis
l. sp. no. 10.
e. v. p. p. p.

Philanthus
1. *Philanthus* *lucivorus* *lucivorus* *lucivorus*
2. *Philanthus* *lucivorus* *lucivorus* *lucivorus*

Philanthus *lucivorus* *lucivorus* *lucivorus*
3. *Philanthus* *lucivorus* *lucivorus* *lucivorus*

Philanthus *lucivorus* *lucivorus* *lucivorus*
4. *Philanthus* *lucivorus* *lucivorus* *lucivorus*
5. *Philanthus* *lucivorus* *lucivorus* *lucivorus*

Philanthus *lucivorus* *lucivorus* *lucivorus*
6. *Philanthus* *lucivorus* *lucivorus* *lucivorus*
7. *Philanthus* *lucivorus* *lucivorus* *lucivorus*

Philanthus *lucivorus* *lucivorus* *lucivorus*
8. *Philanthus* *lucivorus* *lucivorus* *lucivorus*
9. *Philanthus* *lucivorus* *lucivorus* *lucivorus*

Philanthus *lucivorus* *lucivorus* *lucivorus*
10. *Philanthus* *lucivorus* *lucivorus* *lucivorus*
11. *Philanthus* *lucivorus* *lucivorus* *lucivorus*

Philanthus *lucivorus* *lucivorus* *lucivorus*
12. *Philanthus* *lucivorus* *lucivorus* *lucivorus*

Philanthus *lucivorus* *lucivorus* *lucivorus*
13. *Philanthus* *lucivorus* *lucivorus* *lucivorus*
14. *Philanthus* *lucivorus* *lucivorus* *lucivorus*
15. *Philanthus* *lucivorus* *lucivorus* *lucivorus*

Philanthus *lucivorus* *lucivorus* *lucivorus*
16. *Philanthus* *lucivorus* *lucivorus* *lucivorus*
17. *Philanthus* *lucivorus* *lucivorus* *lucivorus*
18. *Philanthus* *lucivorus* *lucivorus* *lucivorus*
19. *Philanthus* *lucivorus* *lucivorus* *lucivorus*
20. *Philanthus* *lucivorus* *lucivorus* *lucivorus*

Philanthus *lucivorus* *lucivorus* *lucivorus*
21. *Philanthus* *lucivorus* *lucivorus* *lucivorus*

Circa lucivorus.

40. *Tenacium* *3ua*
T. fol. subtricuspidatis linear. flor. sessil. lateral. solitar.
in Comitatu Nicæensi & Lincina p[re]q. ad Vias.
Planta morchæm vedolens. Caud. lignos. humilis. hirs.
subtus folia sessilia raro integerrima flor. magnus hirs.
subtus purpureus Variat corolla lutea & alba.
41. — *flavum.*
T. fol. cord. obtuse serrat. bract. integerr. concav. caud.
pubicol. flor. tern. vacuol.
in Agro Nicæensi ad rupes aridas sole p[re]sensitas.
Nicæenses appellant Grosse Germandrê.
Caud. 2. ped. fol. petiol. bas. integerr. undulata dura utrimq.
subnervata supra atro virid. nitid. subt. conciscent. floral.
sessil. concav. Nerviilli multi fl. utrinque 3. pedunculat.
Corol. ochroleuca vel alba raro purpur.
42. — *Capitatum.*
T. capitul. pedunculatis fol. lanceol. crenat. tomentis.
caule erecto.
Minime varium in Comitatu Nicæensi & Liguia.
Habitus majorana sed tomentis. & angustis fl. capitul. sub.
Volunda corol. alb. libur. Violæ.
43. — *polium*
T. capitul. subrotund. fol. ovat. obtus. crenat. tomentis. sessil. caule
prostrato.
in Jaccotis jugi d' Pyrauli. Nicæam versus Cypri.
44. *Lycopus* *variegata*
L. fol. repandis dentatis callosis caud. decumb. corol. cernuis.
in agro Nicæensi.
45. *Cinoglossum cheiripolium*
C. corol. calyc. duplo longior. fol. lanceol. incanis tomentis.
Nicæa crescit Circa Lucerame Cauvini.
46. *Lehim italicum*
L. caud. erecto piloso spicis hirsutis coroll. subæqual. stamin.
longiss.
Locis siccis Nicæensibus Vulgare.
47. — *lulitanicum*
L. caule simplicis fol. caulina. lanceolat. serie. amplissimis
flor. spicat. lateral. coroll. stamin. longior.
in agro Nicæensi.
48. — *plantagineum*
L. fol. radical. ovat. latiss. plantag. fol. simil. lineat petiol.
utrinque ad Vias et loca sicca Nicæa.

49. *Myosotis apula*
in Comitatu Nicaeni.

50. *Orobancha purpurea*
O. caule simplici bracteis tenuis calyce 5-fido.
Locis apricis Satorii Comit. Nicaeni.

51. *Coris monspeliensis* v. pl. Monveg.
in habit. Collet sicco & rupes in agro Nicaeni.

52. *Euphrasia latifolia*
E. fol. dentato-palmatis flor. subcapitatis
in agro Nicaeni.

53. ——— *viscosa*
E. fol. linear calyce glutinose hispido
Orontites fol. viscidis varities serratis.
Locis aridis Collium Nicaenium.

54. *Bartia viscosa*
B. fol. superioribus alternis serratis flor. distant. lateralibus.
in agro Nicaeni.

55. *Bellardia triscago*.
Gen. Calyce bilabiato labio superiore semibifido inferiore
integro Galea vestita barba trifida segmento medio ampliore
Fructus est Capsula ovata biloculari bivalvi polysperma.
Bartia fol. oppositis linear. lanceolatis inciso-serratis spicis
foliorum compactis
Achiranthus calycibus hirsuto-tomentosis fol. oppositis obtuse
serratis caul. simplicissimo.
Nascitur in litore Nicaensi prope Lanterna.
Caul. palmat. fol. 3-uncial. angusta. Spica longa quadrata densa e
cristatis foliis rotundioribus Euphrasiae vulg. simil. compacta.
Fl. magni purpureascentes barbati albi pallentibus Galea vestita
Capsula nigra.

56. *Antirrhinum roseum*.
A. fol. linear. conturb. inferne 4-nis calyc. capit. aequalibus
in agro Nicaeni.
Stavum differt. ab A. monspeliense.

57. ——— *bipunctatum*.
A. fol. linear. glabr. inferius 4-nis caul. erect. paniculata
fl. spicata capitatibus.
in arvis Collium Nicaenium.

59. *Antirrhinum peltarivum*

1. 2. 3. Cant. lineavibus ad radic. oval. sepe tenui-
bus.

t. latioribus

inferiore

dis amplior

hyperma-

strat. spici

uit. obitu

dura e

phacta.

vestita

separatim

maniculat

49. Myosotis apula
in Comitatu Nicæensi.

50

51

52

53

54

55

56

57

58. An

59. An

60

61

62

63

64. Plant

65

66

67

68

69

58. *Antirrhinum peltavianum*
A. fol. ad caul. lineariibus ad radic. ovali saepe ternis
A. fol. caul. altern. radical. lanceol. flor. corymbos.
in agro Nicaensi

59. *Scrophularia auriculata*
S. fol. cordatis oblongis basi appendiculatis subus tormentis
racem. terminalibus.
Ad fontem in agro Nicaensi.

60. ——— *Scorodonia*
S. foliis cordatis duplicato serratis racemo corymboso foliis intermixto
Agri Nicaensis indigena planta est.

61. ——— *betonicefolia*
S. fol. cordat. oblong. dentatis dentibus integerrimis
in agro Nicaensi

62. ——— *peregrina*
S. fol. cordat. lineatis lucid. peduncul. axill. 2. flor. caule 6. angul.
in umbratis Agri Nicaensis.

63. ——— *Lucida*
S. fol. interior. bipinnatis subcarnos. glaberr. racem. 2. parvis.
in agro Nicaensi locis saluberr.

64. *Plantago altissima*
P. fol. lanceol. 5. nerv. dentat. glabr. scap. subangulat. spica
oblonga cylindrica
Jecur le Pailhon prope Nicaeam.

65. ——— *lagopus*
P. foliis lanceolatis subdenticulatis spica ovata hirsuta scap. teret.
in Comit. Nic. copiose

66. ——— *albicans*
P. fol. lanceol. obliq. villis. spic. cylind. erect. scap. teret.
in agro Nicaensi

67. ——— *Bellardi*
P. pubescens fol. sublineariibus plan. petiol. scap. teret. fol.
longitud. spicis cylindricis
invenit. Cl. Bellardi locis sterilibus Villafranca Nica-
ensis prope lanternam

68. ——— *maritima*
P. fol. demicylindricis integerr. basi lanat. scap. teret.
in agro Nicaeae ad rupes.

69. ——— *Coronopus*
frequens in agro Nicaensi.

70. *Plumbago Europrosa.*

Calyx 5. dentatus persistens Coroll. infundibuliformis limbo 5. fido Stamina squamis bacini corollae claudentibus inserta. Stigma 5. fidum Capsula 1. loculari 5. valv. 1. sperma.

Pl. fol. ovato-lanceol. amplexic. ciliat. scabr. fl. congest. Stam. erectiss. in agro Nicaensi.

Caul. 2. cubit purpurascens fol. quandoque obverse ovata atrovirentia florum congesti Calyce erecto tubercul. basi 5. praetunglandul. petiolat. Coroll. purpurascens.

71. *Anagallis Verticillata.*

A. erecta fol. caul. Verticill. axill. folior. in caule & ramis florigeris. flore coruleo. in com. Nicaeni. Cl. Bellard.

72. *Cressa cretica. pentandria Digynia.*

Calyx 5. phyll. persistens. Coroll. tubul. infern. ventricos. limbo 5. fido filamenta tubo insidentia Capsula 1. locut. 2. valv. 1. sperma.

C. coroll. imberbibus capsula 1. sperma.

Nicaeni. agri indigena est prope hospitium di. Villafranca.

Planta parva humifusa incana. caulicul. dodrant. ramis subrubent. fol. minim. altern. sessil. ovat. vel lanceolata fl. inter folia pusilli. sic albo purpurasc.

73. *Juncus Valerandi* Vis. pl. Valder.

ad fontium scaturigines in agro Nicaeni.

74. *Lysimachia linum Stellatum*

L. calyc. coroll. superantibus caule erecto ramorissimo fol. subulatis oppositis peduncul. folio brevioribus. in agro Nicaeni frequent.

75. ——— *tenella.* V. pl. Liguor.

locis aquosis secus flumen le Var & ad oppidum Lucerame.

76. *Gentiana spicata*

G. coroll. 5. fidi infundibuliform. fol. altern. sessil.

in pratis & herbidis locis Nicaeni.

Caulis plerumque unicus subdodrant. fol. opposit. hypervic. mac. fl. termin. albi vel rubelli.

77. *Nerium Oleander*

in montibus Nicae & collibus Lepitellenis.

78. *Perovincia major*

frequent in Comit. Nicaeni.

79. *Aclepias nigra*

A. fol. basi barbati caule superne subvolubili fol. cordat. nudis

flor. confert. patentiissimi petalis nigris.

in collibus Nicaeni & unctis locis Lacois & Edicionibus pariter.

Stem. na
capula 1/2

Stem. erect.

Stem. erect.
capula 1/2

Stem. erect.

Stem. erect.
capula 1/2

Stem. erect.
capula 1/2
fol. 1/2

Stem. erect.
capula 1/2

Stem. erect.
capula 1/2

Stem. erect.
capula 1/2

Stem. erect.
capula 1/2

Stem. erect.
capula 1/2

Stem. erect. flor. fol. cum flor. camp. altitud. plant. paul. ultra 2. 1/2

[Faint, mostly illegible handwritten text in a cursive script, likely Latin, covering the majority of the page.]

flor. confert. parenthemi petal. nigris.
 in collibus & caenibus & unctis locis siccis, & editionibus nascitur.

80. Cym...
 81. Hyos...
 82. —
 83. Verb...
 84. —
 85. Conu...
 86. —
 87. C. p...
 88. C. p...
 89. C. p...
 90. C. p...
 91. C. p...
 92. C. p...
 93. C. p...
 94. C. p...
 95. C. p...
 96. C. p...
 97. C. p...
 98. C. p...
 99. C. p...
 100. C. p...

80. *Cynanchum monspeliacum*.

Gen. Nectario cylindraceo ovifero 5-dentato ab *Asclepiade* differt.
C. caul. volubili herbaceo fol. venis palmi - cordatis acutis

Rarum in agro Nicaeensi nascitur.

Radice & caul. lactiferi fol. opposita longe petiolata integerr. crassa
subalbida fl. axill. fascicul. parvi alb.

81. *Hypocymus albus* V. pl. lig.
in comit. Nicaeensi.

82. ——— aureus V. pl. lig.
ad muros Nicaeae.

83. *Verbascum phlomisoides* V. pl. lig.
in aridiss. & latic. comit. Nicaeensi. abunde Circa Villafanca

84. ——— sinuatum V. pl. ligur.
saepe occurrit in comit. Nicaeensi.

85. *Convolvulus althaeoides* V. pl. ligur.
in comit. Nicaeensi ad super. mari. prociama, & ad muros in vineis.

86. ——— lineatus
C. fol. lanceol. sericeis lineatis petiolat. pedunc. biflor. calyc. sericeis
subfoliaceis.
Nicaeae nascitur.

Caulis prostratus flexuosus semiped. incani fol. remota basi angustiora acuta
ad angulum acutum lineata Hyal. 2. Calyce longiore Corol. dilute
purpurea. erect. hirta.

87. ——— *Cantabrica*
C. fol. lineari-lanceolatis acut. caule erectiuscul. ramoso: pedunc. sub.
2-floris calycibus pilosis.
in agro Nicaeensi abunde

Tota planta usitata caul. sesquiped. fol. altern. integerr. radical. petiolat.
obtusiora; caulina. serrit. pedunc. axill. cor. carnea c. lineis rubris.

88. *Campanula medium* C.
capitul. 5-locul. obtechi; caule indiviso erecto
glossoso floribus erectis.
in Sylv. / apricis & locis incultis agri Nicaeensis.

89. ——— *erinus*
C. caul. dichotomo fol. serrit. superior. oppos. bidentatis
agri Nicaeensis indigena.

90. *Colchicum montanum*.
C. flore cum foliis conjuncto petal. lineari.
in praet. macilentis montanis non tamen editioribus Circa Cospeho.
altiss. mont. flor. fol. cum flor. erump. albis plant. paull. ultra 2. g.

91. *Agave americana*. *Hernand. monogyn.*
 Gen. coroll. erecta superba & spida epicaupia. Stamina corolla lon-
 giora capsula 3. locul. 3. valv. polypetala. fol. Carnos. succulent.
 A. fol. dentato spinos. scap. amolo.
 Ad vases in agro Nicaeensi inter Villafranca & Monaco.
 Fol. applanatissima succulenta hinc convexa inde plana mucrone durissimo
 firmi. peroram spinis obita. Scap. erect. etiam 30. pedat. simplici
 spica ramosa fl. innumeri petala sublatea.

92. *Erica scoparia* V. sp. Lig.
 C. Bellardi in montibus di. Cathon in Comitatu Nicaeensi.

93. *Cotyledon umbilicus* Vener. V. sp. Ligur.
 in Comit. Nicaeensi. ad muros & vases

94. *Zarminum fruticosum*
 ad sepes in agro Nicaeensi

95. ——— officinale
 frequent in comitatu Nicaeensi

96. *Olea europaea*
 in Comitatu Nicaeensi

97. *Phillyrea media* *Dianthia monogyn. L.*
 Calyx 4. dent. corolla 4. fida rotata Baccæ monosperma.
 sp. fol. ovato-lanceolatis subintegerrimis
 locis sicis & vasis in agro Nicaeensi.

98. ——— angustifolia
 sp. fol. linear. lanceol. integerr.
 iidem crescit locis

99. ——— latifolia
 sp. fol. ovato cordatis serratis.
 Orietur ad vases Montalbani in agro Nicaeensi.

100. *Momordica elaterium*.
 in agro Nicaeensi provenit.

101. *Nitescens agnus cathus* V. sp. Ligur.
 inter Mentan. & Monaco.

102. *Lycium europaeum* *Pentandria monogyn.*
 Cal. 5. dentatus coroll. in fundibuliform. Stam non defluunt. Bacc.
 villis. baccæ 2. locul. polypetala.
 fol. obliq. ramul. flexuos. teret.
 Fructus ramulosi. fol. 5. 5. etc. ead. gemma prodeunt fl. e singulis gemmis
 2. 5. cal. 5. fida coroll. fauce pallid. nigro. int. int. purpurea.

Colaba (on)
... ..

... ..
... ..
... ..

... ..

... ..
... ..

... ..

... ..

... ..

... ..

... ..
... ..

91. *Agave americana*. *Hesperid. min.*

92

93

94

95

96

97

98

99

100

101

102

Frut. c. 1 m. alt. fol. 5. 6. etc. ead. gemma prodeunt fl. e. lingul. gemm.
2. 5. cal. 5. fidus corol. fauce pallid. nigro. int. purpureo.

103. *Viburnum lincx*
V. fol. ~~integerrimis~~ ovatis ramificationibus Ucnarum subny
Villato glandulosis.
in Sylvis opacis Nicaeensis.

104. *Thymactea laevicola*
Daphne racemis axillaribus 5. flor. fol. lanceol. glabris
Tota planta flexilis supra medium ramosa ramis apice folios pubescent. 2. pedat.
evidens cinereo-fuscus. Fol. digit. longa laminae firmitate alterna breviter
petiol. perennant. crass. integerr. supra laev. virid. subt. dilutiora.
Racemi nutantes bracteis concavis alternis Fl. frutres.
in Sylvis umbris non humidis Mont. Comit. Nicaeensis. palium.

105. *L. lanamunda.*
D. thymactea flor. sessil. axillar. fol. lanceol. caul. simpliciss.
Caul. plant. palm. fl. sparii non racemosi corol. et alba hutea.
Circa Nicaeam.

106. *tarcon-raila.*
D. fl. sessil. aggregatis axillar. fol. ovatis utrinque pubesc. nervos.
in agro Nicaeensi.
Futiculus fol. mollia candicant. & velut. Sericea fl. albo pall.

107. *quidium*
D. panicula terminali fol. linear-lanceol. -acuminat.
Frequens in Comitatu Nicaeensi.
Caulis humilis. fruticosa. Cubit. & ultra ramos. fol. Viridis. annua glabra.
fl. congest. alb. supra rubri cinct. fol. suaveolent. fruct. matur. rubri
succosi Bis quatuor annis florent.

108. *Arbutus Unedo* Decandr. Monogyn. h.
Cal. brev. 5. fid. petiol. Coroll. campanuliform. globos. 5. fid. Ovar.
intra florem. fructus 5. locut. polysperm.
A. caule arboreo fol. glabr. serratis baccis polyspermis.
in agro Nicaeensi abunde. Arbor.
Frutere magnas. fol. laevissim. altern. petiol. oblong. duras late
viridia nitida corol. herbido rubra Balca magna sphaerica. rubi-
cunda hirsuta. sem. exasperata.

109. *Styracis officinale.* Decandr. monogyn. Cal. 5. dent. hypocrater.
Corol. imbriculisperm. in 5. vel 6. profund. lacin. divisa fructus est
dupla 1. locut. 2. sperma.
In agro Nicaeensi. abunde in rup. marit. prope Joet.
Arbut. fol. altern. petiol. oval. supra virid. nitent. subt. toment. mal.
Coton. simil. sed min. pedunc. saepe 2. flor. Cor. alba.

110. *Scabiosa leucantha*
S. corollul. 4. fid. subgynat. squam. calyc. ovat. obtus. fol. pinnatifid.
in collibus aridis Nicaeensi. abunde.

111. — *integrifolia*
S. corollul. 4. fid. fol. indivis. radic. ovat. sess. ramis lanceol. caul. herb.
abunde secus arva in agro Nicaeensi.

112. — *grammifolia*
S. corollul. 5. fid. calyc. breviss. fol. caul. bipinnat. filiformibus.
Nicaeensis agri indigena est. *Scabiosa stellata*. addenda.

113. *Globularia alyssum*. V. pl. figur.
in sylvis et incultis agri Nicaeensis.

114. *Ambrosia maritima*. Monoclea pentandria.
chalc. cal. comm. 1. phyll. 5-8 fid. corol. inmundibul. 3. fid.
recept. nud.

Form. cal. 2. phyll. integ. ventr. 5. dentat. 1. flor. corol. o. nuda
e calyce indurata monosperma.

A. foliis multifidis spicis solitariis pilosis sessilibus.
in arenosis maritimis Nicaeensibus.

Caul. subped. ramosis puber. sicut fol. & racemi folia saepe
opposita petiolata mollissima albescentia Racemi densi fl. subrecti.
conferti corol. vivid. lutea.

115. *Xanthium spinosum* V. pl. figur.
ad maris latus occurrit.

116. *Echinops ritro*.

E. foliis supra glaberrimis subtus tomentosis pinnatifid. capitul.
globos. caul. multiflor. corymbos. flor. caeruleis.
abunde in montibus Nicaeensibus.

117. *Carduus leucographus*

C. fol. decurrentibus dentatis spinosis peduncul. nud.
longissimis uniflor. calycibus spinosis inclinator.
Citr. maculis argenteis notat.
in olivis & rudibus circa Nicaeam.

118. *Cirsium oleraceum*

Cirsium fol. pinnatif. farinatis nud. amplexic. inerm. bracteis
subcoloratis integris concavis flore flavicante
in humoribus Nicaeensis agri.

110.

111.

112.

113

114

11

11

1

119.

120.

121.

122.

123.

124.

125.

126.

127.

128.

129.

130.

131.

132.

133.

134.

119. *Cirsium tuberosum*.
C. radicibus tuberosis fol. semipinn. pinn. angulos. ciliat. fl. nud.
Ubique in agro Nicaensi.

120. — *stellatum*.
Carduus fol. integr. sessil. inerm. subt. tomentos. pinn. ramos.
aristat. flor. sessil. lateral.
in agro Nicaensi ad viam securaria.

121. *Acarna cancellata*. *Attractyl. Linn.*
Gen. Calyce circumdat. bract. foliac. corollat. hermaphrod. 5. fid.
receptac. paleac. paleis in 2. aut 3. aristas abeuntibus in nostra
planta sem. Villois. papp. plumos. coronat.
Attract. involucri cancellati. Ventr. linearibus dentatis ca-
lycis ovatis floribus flosculis
ad Rupes Montalbani in agro Nicaensi et ejusdem Comit.
loc. steris. praecalis.

122. *Carthamus tinctorius*.
Ubique in ar. di. collibus Nicaensibus.

123. *Carduncellus mompeliensis*. *Carthamus L.*
Gen. Calyce imbricatus squamis appendice foliacea auctis floscul.
omn. hermaphrod. limbo 5. fido receptac. paleac. sem. papp. simpl.
coron. Carth. fol. contin. linear. pinnat. longitud. plantae.
in montibus interpositis oppido Lucevame & Lantolen.

124. *Cnicus ferox*. V. M. Tard.
Cl. Dana in collibus di. Braun.

125. — *acarna*
C. fol. decurr. lanceol. indivis. calyc. pinnat. pinnol.
in Comit. Nicaens. freq. loc. steris. siccis. & ad vias.

126. *Carlina lanata*
C. caud. bipido calyc. sanguin. terminal. primo aristati sessil.
ad maris litorum locis siccis.

127. — *achanthifolia*
C. acut. fol. tomentos. semipinn. plicatis pinn. semipilob. pinn.
gentibus.
In montibus Braun alique soppella praecians. frequens.

— Cent. panicul. Var. flor. albo ad rupes in agro Nicaensi.

128. *Centauria cineraria*
C. calyc. ciliat. terminali sessilibus fol. tomentos. bipinn.
bipidis lobis 7 acutis.
in montibus circa Lucevame.

129. *Centaurea sonchifolia*.
C. calyc. palmatis spinosis fol. decurrentibus spinulosis
veppando-dentatis
loc. 7 aridis collium Nicæensium.

130. ——— *seridis*
C. calyc. palmatis spinosis fol. decur. toment. orbicongis inflm.
sinuato-dentatis.
in agro Nicæensi

131. ——— *ilwardi*
C. calyc. palmatis spinosis fol. lyvato-dentatis hispidi
subamplexicaulibus flor. sess. f. feracinal.
in vineis Nicæensibus & avenis maritimis.

132. ——— *aspera*
C. calyc. palmatis spinosis fol. lanceolat. dentat.
locis sterib. Comit. Nicæens.

133. ——— *Nicæensis*
C. calycibus ciliatis spinosis fol. virid. asper. ovat. radi-
cal. petiol. dentat. caul. n. amplexicaul. decurrent.
A. Bellardi circa Nicæam legit.

134. ——— *galactites* V. pl. Ligur.
ubique abunde in Comit. Nicæens.

135. *Tanacetum annuum*
T. fol. bipinnatifid. linear. acut. corymbis tomentis.
Nicææ, et in arvis Nicæens. Sicut flumen le Varanon. in frequent.

136. *Anacyclus Valentini*
Gen. Calyx imbricat. hermiphor. placent. paleac. flor. disc. 5-fid.
hermaphrod. sem. compress. nuda in amb. h. form. paucae revol.
lingul. brev. integerr. sem. alata apic. emarginata.
A. fol. decompositis linearibus lacin. div. feretiuscul. acut.
flor. flosculot.
in Agro Nicæensi

137. *Artemisia Gerulescens*
A. fol. caul. lanceol. integr. tomentos. radical mul-
tipis. flor. long. spicat. tenuis.
Nicæense litus inhabit.

Herbertus...
...
...
...
...

Herbertus...
...
...
...
...

Herbertus...
...
...
...

Herbertus...
...
...
...

Herbertus...
...
...
...
...
...
...

Herbertus...
...
...
...

Herbertus...
...
...

Herbertus...
...
...

138. *Microglossa*
 Calyc. 2
 5. 2
 139. *Filago*
 F. flor.
 in
 140. *Guaiacum*
 141. *Stachys*
 Calyc. 2
 142. *Stachys*
 Calyc. 2
 143. *Corymbosus*
 C. fo.
 acu.
 144. *Asperula*
 145. *Asperula*
 146. *Asperula*
 147. *Asperula*
 148. *Asperula*
 149. *Asperula*
 150. *Asperula*
 151. *Asperula*
 152. *Asperula*
 153. *Asperula*
 154. *Asperula*
 155. *Asperula*
 156. *Asperula*
 157. *Asperula*
 158. *Asperula*
 159. *Asperula*
 160. *Asperula*
 161. *Asperula*
 162. *Asperula*
 163. *Asperula*
 164. *Asperula*
 165. *Asperula*
 166. *Asperula*
 167. *Asperula*
 168. *Asperula*
 169. *Asperula*
 170. *Asperula*
 171. *Asperula*
 172. *Asperula*
 173. *Asperula*
 174. *Asperula*
 175. *Asperula*
 176. *Asperula*
 177. *Asperula*
 178. *Asperula*
 179. *Asperula*
 180. *Asperula*
 181. *Asperula*
 182. *Asperula*
 183. *Asperula*
 184. *Asperula*
 185. *Asperula*
 186. *Asperula*
 187. *Asperula*
 188. *Asperula*
 189. *Asperula*
 190. *Asperula*
 191. *Asperula*
 192. *Asperula*
 193. *Asperula*
 194. *Asperula*
 195. *Asperula*
 196. *Asperula*
 197. *Asperula*
 198. *Asperula*
 199. *Asperula*
 200. *Asperula*

138. *Microglossa*
 Calyc. 2
 5. 2
 139. *Filago*
 F. flor.
 in
 140. *Guaiacum*
 141. *Stachys*
 Calyc. 2
 142. *Stachys*
 Calyc. 2
 143. *Corymbosus*
 C. fo.
 acu.
 144. *Asperula*
 145. *Asperula*
 146. *Asperula*
 147. *Asperula*
 148. *Asperula*
 149. *Asperula*
 150. *Asperula*
 151. *Asperula*
 152. *Asperula*
 153. *Asperula*
 154. *Asperula*
 155. *Asperula*
 156. *Asperula*
 157. *Asperula*
 158. *Asperula*
 159. *Asperula*
 160. *Asperula*
 161. *Asperula*
 162. *Asperula*
 163. *Asperula*
 164. *Asperula*
 165. *Asperula*
 166. *Asperula*
 167. *Asperula*
 168. *Asperula*
 169. *Asperula*
 170. *Asperula*
 171. *Asperula*
 172. *Asperula*
 173. *Asperula*
 174. *Asperula*
 175. *Asperula*
 176. *Asperula*
 177. *Asperula*
 178. *Asperula*
 179. *Asperula*
 180. *Asperula*
 181. *Asperula*
 182. *Asperula*
 183. *Asperula*
 184. *Asperula*
 185. *Asperula*
 186. *Asperula*
 187. *Asperula*
 188. *Asperula*
 189. *Asperula*
 190. *Asperula*
 191. *Asperula*
 192. *Asperula*
 193. *Asperula*
 194. *Asperula*
 195. *Asperula*
 196. *Asperula*
 197. *Asperula*
 198. *Asperula*
 199. *Asperula*
 200. *Asperula*

138. *Micropus supinus*.

Calyx. 5-phyll. calyculatus placenta nuda flor. discipnati corol.
5. dentata styli rudim. habentes & abortientes in ambitu flores
feminei nudi sem. nuda squam. Calyc. inclusa.
Chr. caul. procumb. fol. gemin. rachibus muricatis
Nicaea ad maris litus.

139. *Filago acaulis*.

foliis
F. flor. acaulis sessil. floral. majoribus
in stagnis exsiccatis Nicaeensis

Raro caulescit id accidit locis fertilibus & tunc. caul. superius sa-
moli. ram. prostr. inforam. fol. imbricata linearia tomentosa - in-
cana. Fl. alii majores alii minores Nicaeensis.

140. *Guaphalium Stachas*.

G. pubesc. fol. linear. ram. virgat. cymis. compositis.
in aridis collibus Nicaeensis abunde.

141.

luteo album

G. herbaceum fol. semiamplexice. ensiform. repand. obtus.
utrinque pubescent. flor. conglomerat.

Abundante cum rivulis & fluentes aquas in agro Nicaensi.

142. *Stachelina Dubia* L.

Calyx. oblongus imbricatus ex squamis coloratis flor. herma-
phrod. 5. fil. placenta paleacea semina. papp. plumos.

J. fol. linear. denticulatis squamis calycinis lanceolatis
pappo calyc. duplo longiore.

in Agro Nicaensi.

Caul. ped- cubit incano-tomentosi. rami pl. erecti circ. med. saepe
iter. ramosi. fl. 2-3. conglomeratis fol. plur. inordinat. sessil. fl. oblongi.

Calyx. squamulis purpurascens

143. *Corypha Sarcobatis*

C. fol. linear. subdent. peduncul. long. 7. uniflor. capitulo
acuto & angust.

frequent. ad muros in Comit. Nicaensi.

144. ———— *Sordida*. V. pl. Lign.

ad Rupes & Vetus muros in Com. Nicaensi.

145. *Buphtalum*. Lignos. V. pl. Lign.

in lignis ad Rupes & vias in agro Nicaensi.

146. *Buphtalmum agrarium*
B. calyc. obtuse foliis. Jovis. arell. fol. altern. oblong. obtus. caul. hirsut.
 in agro Nicaeensi: ad font. Scaturigines.
147. ——— *maritimum*
B. calyc. obtuse foliis. peduncul. fol. altern. spatul. caul. hirsut.
 in agro Nicaeensi: prope marit. litus.
148. *Achillea ageratum*
A. fol. lanceol. obtus. acut. serrat. flor. umbell. latis.
 in Comit. Nicaeensi. Frequens circa la Turbia et secus flum.
 spallion in arenosis.
149. *Chamaemelum maritimum*
Anthem. fol. pinnat. dentat. Carnos. nud. & pinn.
 stat. caul. prostrato calyc. sub tomentol.
 in agro Nicaeensi.
150. ——— *tomentosum*
A. fol. pinnatif. obtus. plan. pedunc. hirsut. foliis. calyc.
 tomentol.
 in arenosis maritimis agri Nicaeensis.
151. ——— *mixtum*
A. fol. simplic. dentat. laciniat. coronop. fenis.
 in Coll. Nicaeensi. mar. proscium.
152. ——— *nobile*
A. fol. lanuginos. pinnat-composit. linear. acut.
 in Comitatu Nicaeensi.
153. ——— *valentinum*
A. caul. ramor. fol. pubesc. 3. pinnat. setac. calyc. villos.
 peduncul.
 in tabulor. flum. spallion in agro Nicaeensi.
154. *Calendula arvensis*
 Jecur arva in agro Nicaeensi.
155. *Cysanthemum discordium*
Cohula fol. altern. lanceol. dentat. caul. erect. fl. solit.
 ubique locis aridis in Comit. Nicaeensi.
156. ——— *inodorum*
C. fol. pinnat. multifid. caule ramoso diffuso.
 inter segetes in agro Nicaeensi.
157. ——— *montanum*
C. fol. imis spatulat. lanceol. serrat. summi. linear.
 in monte Syran aridis & salscoris locis uatior.
 Vix a *C. leuchanthemo* differt.

47.

4.

1874.

24

гит в.

in monhous circa 1916 same Cl. - 1/2 Edward prop. 1/2 monhous circ.

146.

147.

148

149

150

151

152

153

154

155

156

Vix a C. leucanthemo differt.

longum

647.

sup.

plum.

pl.

orgumb.

in montibus circa Lucerne Cl. Helveti proprietatem C.C.

146

:

147

148

149

150

151

1

152

153

154

1

Vix a C. Leuchanthemo differt.

155 Chrysanth

156

157

Cant. 2. h.
et integr.
fari pale

158 Belli
A.
in

159 Astro

C.

160

161

162 Evig

163

164 Senecio

165 Ciner
Abunde a
supra

166 Lappan

Lappan
in me

158. *Chrysanthemum Segetum*
C. fol. amplexic. superm. lac. in. inferm. dentato-serratis.
in salubrior fl. & Var. et ad marit. litus.
159. ——— *myconii*
C. fol. lingulat. obtus. serrat. Calyc. squam. aequalibus.
in agro Nicaeensi.
160. ——— *coronarium*
C. fol. pinnat. incis. echinorum latioribus
Cum superiori repperit. est a D. Grudice.
Caul. 2. ft. pedat. Ramor. sulcat. folios. fol. sessil. pinnisveb. semipinn.
vel integr. pinnat. Echrem. major. 2. fid. moll. virent. flor. ampl.
lutei pallescentes.
161. *Bellis annua*
fl. minima praeterea caule folioso.
in gl. h. g. v. minoribus Nicaeensibus sed varia.
162. *Aster Hippolium*
C. fol. lanceol. integerr. Carnor. glabr. Ram. inaequat. flor. corymb.
ad mar. Nicaeensi. litus loc. paludos.
Caul. cubit. Ramor. 1. triat. Ruffere. pedunc. nud. 1. flor. Calyc.
squam. obtus. corol. rad. corul. Discus luteus.
163. ——— *acris* V. pl. liguv.
in Comit. Nicaeensi.
164. ——— *bifrons* V. pl. liguv.
in Com. Nicaeensi. non in freq.
165. *Erigeron Viteorum*
C. peduncul. 1. flor. lateral. fol. lanceol. dentat. Gali refle-
xer. Calyc. squam. coroll. radiat.
in Comit. Nicaeensi.
166. ——— *graveolens* V. pl. liguv.
loc. 7 audit agr. Nicaeensi.
167. *Senecio Sylvaticus* V. pl. liguv.
abunde in C. Nicaeensi.
168. *Cineraria maritima* V. pl. liguv.
Abunde ad marit. Nicaeensi. litus nec non ad v. sup. Roque tagline
supra l'elcarene.
169. *Lappana fatida* Calyc. ex squam. parv. alt. unius ordinis. Sem.
nudum.
Lappana caule nudo indiviso fol. semipinnat. pinn. retraversi. dentat.
in montibus circa Lucerame et Bellard prope Carthou Nic.

170. *Lappula stellata*

l. calyc. fructus undiq. patentibus rad. subul. fol. caulic. lanceol.
indivisi.
Nicæa observata est.

171. *Catananche cerulea* V. M. figur.

abund. in sterilibus Comit. Nicæens.

172. — *lutea*

C. squam. calycin. inferior. lanceol.
prope Breglio.

173. *Scotymus maculatus*.

Calyc. imbric. sem. coronat. avit. 2. aut 3. facile decidui.
J. floribus solitariis
Anelice & Nicæa ad mare litus.

174. — *hispanicus*.

J. floribus congestis.
in agro Nicæensi.

175. *Wewia barbata*.

Cal. ad basin calycul. ordine squam. linear. sem. tam disci quam
peripherie avit. 2. instruct.
Crepit involucri. Calyc. longior squam. setac. spuri.
in agro Nicæensi ad limites agrorum.

176. *Leitodon bulbosum*

l. fol. oblong. ovat. subdent. glabr. calyc. lœv. scap. super.
hirsut.
Nicæensi & agri indigena planta est.

177. *Picris tuberosa* V. M. figur.

in privat. Nicæensibus frequens est.

178. *Hieracium Capillaceum*

H. caul. tubund. uniflor. fol. nervos. glavin. integerr.
hectum circa Soppello.

179. *Crepis burrifolia*

C. fol. pinnatifidis crenatis scapo nudo panicifloro.
Agri Nicæensi indigena planta est.

Page 112

cauliculus

in: 112

in: 112

11

in: 112

integre

[Faint handwritten text, likely bleed-through from the reverse side of the page. The text is mostly illegible due to fading.]

Urtica dioica L.
Large perennial herb, leaves serrated, stinging hairs.
Common in wet places.

Urtica dioica L. var. *diandra* (L.) Kunt.
Similar to the above but with different leaf shape.

Urtica dioica L. var. *diandra* (L.) Kunt.
Similar to the above but with different leaf shape.

Urtica dioica L. var. *diandra* (L.) Kunt.
Similar to the above but with different leaf shape.

Urtica dioica L. var. *diandra* (L.) Kunt.
Similar to the above but with different leaf shape.

Urtica dioica L. var. *diandra* (L.) Kunt.
Similar to the above but with different leaf shape.

Urtica dioica L. var. *diandra* (L.) Kunt.
Similar to the above but with different leaf shape.

Urtica dioica L. var. *diandra* (L.) Kunt.
Similar to the above but with different leaf shape.

Urtica dioica L. var. *diandra* (L.) Kunt.
Similar to the above but with different leaf shape.

Urtica dioica L. var. *diandra* (L.) Kunt.
Similar to the above but with different leaf shape.

180. *Crepis albidus*
C. fol. duncinato-dent. incan. squam. calyc. simpl. membranac.
fl. albid.
 Com. Nic. Juncos montani redibit. apparet in Com. Nic. aene.
181. — *vulva*
C. fol. amplexic. lyvat. duncinat. f. h. fl. suave rubenti.
 in Com. nic. secus arva inter Soppello & Scarenese
182. — *putchva* V. *fol. ligur.*
 in Vineis & oliveis Nic. aene.
183. — *nemantensis*
Hieracium sanctum Jacq. nud. multiflor. fol. lyvat. obtus. dentat.
 in sterilibus circa Quatara in Com. Nic. aene.
184. — *pyrengeiana*
H. caul. varior. folior. fol. semiamplexicaul. oblong. repand. hisp.
 ad limites agrorum in Comit. Nic. aene.
185. — *echioides*
Picris perianthii exterior. 5. phyll. inter. avit. to majoribus
Regius in agro Nic. aene.
186. *Louchus maritimus*
L. fol. omn. integr. denticul. spinos. ram. 1. flor. semiflorul.
5. dentat.
L. peduncul. nud. fol. lanceol. amplexic. indivis. rebos. argut.
dentat.
 in humidis & ad fluent. ag. in agro Nic. aene. nascitur.
187. — *picroides*
Scorzonera fol. superior. amplexicaul. integer. infer. duncin. pe-
dunc. squamos.
 in Com. Nic. aene.
188. *Lactuca saligna*
L. fol. im. semipinn. super. amplexic. hamat. nerv. spinos.
 in Com. Nic. secus arva copiose evasit.
189. *Prenanthes viminea* V. St. Jend.
 in Com. Nic. prope le Scarenese Soppello.
190. — *variorissima*
Lactuca varior. pene ligur. fol. radical. pinnatif. caul. lanceol.
 in salubris ad maris Nic. aene. libet.

191. *Rhagadiolus hedyotidis*. Hyoseris vel Lappiana L.
Gen. demum a radic. Sappius Nuda Gen. Valia Sappio aut Siquani,
Coronata.
Hyoser. fruct. subglobosis glabris caule ramoso
in vineis et arvis incultis agri Nicæensis nascitur et in arenis.
Semen fl. Sappion.
192. ——— creticus
H. fruct. ovatis scabris caule ramoso.
Semen arva in agro Nicæensi.
193. ——— scaber
H. scapp. simplicissimo l. floro fol. scabris. Sem. Sappios.
in olivet. Villa franca montis alban: ad mania sub-elliptica.
194. ——— *Racineka*
Lappiana calyc. fruct. torulosis depress. obtus. sessil.
loca inculta mari Nicæensi: proxima inhabit.
195. ——— radiatus
H. scapp. uniflor. nud. fol. glabris runcinatis angulis dentatis.
in collibus maritimis Nicæensibus lectus est.
196. ——— *karavencoides*
H. fol. sinuato-dentatis Sem. central. Sappio. Plumos. pilis folior.
juventibus radice fibrosa
in Rupibus circa Lucernam legit D. Cavvini.
197. *Scorzonera hispanica*
J. caule ramoso fol. amplexic. integ. serrulatis.
in pratis comit. Nicæens.
198. ——— *laciniata* N. Pl. Vall. Macra.
in com. Nicæens. freq. est.
199. *Tragopogon Dalechampii*
J. calyc. l. phyllis. corol. brevior. inerm. fol. runcinatis.
in vineis & arvis. Nicæens.
200. ——— *picroides* V. Pl. Liguor.
in Olivetis Nicæensibus.

piculat.

191

19

193

194

195

196

197

198

199

200

201. G

202. T

203. H

204. F

205. A

207. Cn

208. Tr

209. Ba

201. *Geropogon glabrum*.
Cal. 4-phyll. profund. 8-fid. Sem. duci pappi plumos. Rad. arit. denticulat.
coronata.
9-fol. glabrii graminei; flore dilute incarnato.
in agro Nicaeni.

202. *hirsutum*.
9-fol. pilosis caule hirs. calyc. corol. brevior. peduncul.
superne inestallat? fl. luteo.
in herbidiis collibus Nicaenibus.

203. *Hypochaeris radicata*
H. fol. vuncinatis obtusis scabris caule ramoso undulato
pedunc. squamul.
in agro Nicaeni; secus arva frequens.

204. *Seriola urens*. Gen. Calyc. excl. ord. squamarum. Sem.
omnium pappi plumosus.
J. caule ramoso foliolo fol. dentat. calyc. hirsid.
ad vias et loca sterilia Comit. Nicaeni.

205. *Andriata integrifolia*
Calyc. ex unico ordine squamarum pappi simplici.
A. fol. inferior. vuncinat. superior. ovali. oblong. tomentis.
in sterilibus agris Nicaeni.

206. *sinuata*.
A. fol. vuncinatis
Secus flumen le Var in agro Nicaeni.
Cl. Vilmorin ad precedentem refert.

207. *Cneorum tricoccon*.
Gen. Calyc. 3-dent. persistens petal. 3. aequal. Stam. 3. Stigma
3-fid. Capsula tricoeca.
Rupes agri Nicaeni inhabitat.
Fruticulis fol. serrul. oblong. angust. integerrima peduncul.
olitar. 1. flor. corol. lutea Bacca rubra dein fusca

208. *Triglochin maritimum*.
T. capsul. 6-locularibus ovalis
locis palustribus maritimis mari proximis.

209. *Batonia tenuifolia* Tetrandria Digyn.
Calyc. 4-phyll. persistens foliol. altera brevior. petala 4. calyce
breviora angusta stigmata 2. Capsula 1-locul. 2-valv. 2-sperma.
in pascuis sterilibus Comit. Nicaeni.

210. *Alytrum maritimum* V. *Pl. Ligurica*.
interstitibus agri *Nicaeensis* mari proximi.
211. — *Sativum*
Vern. Ravillon *Iponte* in agro *Nicaeensi*.
212. *Junaria Clypeata*
Alytrum caule erecto herbaceo *Silicul.* *Settis.* oval. compressis.
plant. petal. acuminat. linear.
in agro *Nicaeensi*.
213. — *halymifolia* V. *Pl. Vall. Macra*.
frequens in Comit. *Nicaeensi* prope *Sorpetto*.
214. *Fosselinia jonthlippi* *Silicula* 1. locular. orbicul. plan. 1. *hypem.*
Clypeola *Silicul.* orbicul. 1. locular. 1. *hypem.*
in agro *Nicaeensi*.
caulicul. plant. sub palm. ex rubore obscuro incana fol. *Serpilli* altern.
cylindrica incana fl. terminat. racemosi parvi petal. luteola.
215. *Biscutella auriculata* Vid. *Pl. Tenu*.
inter seget. Comit. *Nicaeensi* circa *Sorpetto*.
216. *Thlaspi hirtum*
J. *Silicul.* sub. tundi pilosis fol. caulic. sagittat. villis.
in *Nicaeensi* & *Nicae*.
217. *3. beris sempervirens*
3. frutescent fol. linear acut. integerr. fl. albo.
in sabulosis flum. *Spallien* *Nicae*.
218. — *umbellata* A
3 herbacea fol. acuminat. inferior. serrat. superior integerr.
in Comit. *Nicaeensi*.
219. *Lepidium procumbens*
l. fol. pinnatifid. extremo major. scap. nud. caul. prostr. racemifer.
in aridit agri *Nicaeensis* videtur.
220. — *spetueum*.
l. fol. pinnat. integerr. petal. emargin. calyc. minor.
ad super locis aridis in Comit. *Nicaeensi* abunde.
221. *Cochlearia officinalis*
in *Spallien* *lappidosis* alpinis di. *Santoca* *Iponte* nascitur.
222. — *diva* A
C. fol. lanceol. dent. amplexic. incana.
loc. incult. & sec. arv. in agro *Nicaeensi*.

in agro Nicaeni prope Villanica.

210. *Alyrium maritimum* V. M. L. L. L. L.

211.

212.

213.

214.

215.

216.

217.

218.

219.

220.

221.

222.

23. *Cram.*
Ho.
1-6.
Caul.
Petal.

34. *Linop.*

35. —

36. *Bra.*

37. —

38. —

39. —

40. *Hyp.*

41. —

42. *Che.*

43. —

44. —

45. —

223. *Crambe maritima*
 Fruct. subglob. non delitens deciduus baccaam siccam interdum reperiunt
 1-locularis aut 2-locular. sed. 1-sperm. alter. locul. abortiente.
 Caut. 3. pedat. fol. magna sinuato-pinnatifid. carnos. succulent. glauca.
 petal. majorecula alba.
 ad Libus Nicaeensis.
234. *Sinapis alba.*
 Siliq. hispida. tort. obliq. longit. euriform. sem. albo & ruffo.
 loci pinguius in agro Nicaeensi.
235. — *maritima* V. pl. Tend.
 in montibus Versus Nicaeam.
236. *Brassica orientalis*
 B. fol. cord. amplexic. glob. radical. scabr. integerr. Siliq. 4-goni.
 in Cons. Nicaeensi. Lecta est flor. albam proferit.
237. — *arvensis* ^{amplexicaul. cordatis}
 B. fol. spatul. repandis summ. ~~linear. lanceol.~~ integerrimis
 B. campestris pinnatifida fl. purpureo foliis pinnatis.
 Ad vias & muros hortorum circa Ventimiglia.
238. — *alpina*
 B. fol. cantin. cord-sagittatis amplexic. radic. ovat. petal. erectis.
 locis umbratis & abruptis Cons. Nicaeensis circa la Giandola
239. — *eruca*
 Nicaea in incultis reperitur aruga Vern. nobis
240. *Heperis verua*
 H. caul. erect. ramos. fol. cord. amplexicaul. serrat. villor.
 Habitat in agro Nicaeensi
241. — *laciniata* Vid. pl. Tend.
 Circa Soppello.
242. *Cheiranthus Borone*
 inter Villafranca & Menton. locis aridissimis.
243. — *maritimus*
 C. fol. lanceol. acutiusculis caul. ramos. diffus. anther. emittunt.
 Habit. in agro Nicaeensi.
244. — *litoreus*
 Ch. fol. lanceol. subdentat. subfomentol. subcarnos. petal. emar-
 ginat. siliq. tomentol.
 Hae litoreis Nicaeensis indigena planta est
245. — *sinuatus*
 C. fol. tomentol. obtus. subinnat. ram. integerr. Siliq. muricat.
 in agro Nicaeensi prope Villafranca.

246. *Oxylinum cheiranthoides*.
O. fol. lanceol. integerrimis, siliquis parvis
 nascitur in agro Nicæensi.
247. *Silymbrium murale*
S. subcaul. fol. lanceol. sinuato serrat. laeviusculis, scapis subulatis
 adscendentibus.
 Secus arva frequens in agro Nicæensi.
248. — *Vimineum*
S. acaul. fol. lyrat. laev. scapp. adscendentibus fl. minut.
 Circa Joppitellum Onoratus Causini legit.
249. — *Barbellei*
S. caul. subnud. ramor. fol. radical. lanceol. dentat. hispida.
 Secus arva in agro Nicæensi.
250. *Ruta graveolens*
 in comit. Nicæensi.
251. *Euphorbia chamaerice*
E. dichotom. fol. crenulat. subrotund. glabr. fl. solit. axillat.
 caul. prostrum.
 Ubique locis aridis agri Nicæensis.
252. — *peplis*
E. dichot. fol. integerr. semicordat. flor. solit. axill. caul. prostrum.
 ad litus maris Nicæensis.
253. — *escigua*
E. umbella 3-fida dichot. involucl. lanceolat. fol. linear.
 in agro Nicæensi frequens.
254. — *spinosa* V. pl. figur.
 Ubique ad Joppitellum in com. Nicæensi.
255. — *epithymoides* V. pl. figur.
 locis incultis & ad oras sylvarum com. Nicæensi.
256. — *Nicæensis*
E. glabra caul. subsimplici umbella 5-fida 2-fida fol. ovat.
 capsul. laevibus non sulcatis
 in com. Nicæensi inter Cimie & la Trinità.
257. — *navalis*
E. umbella sub-5-fida bipida involucl. cordat. reniformibus
 fol. surr. imbricat.
 ad maris Nicæensis litus & in sabulosis.

long. 1/2

ant.

long.

lat. 1/2

ant. 1/2

linear.

st.

foliis

la.

reniformis

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

218. *Agrostis* *peruviana*
L. *peruviana* *peruviana* *peruviana*
L. *peruviana* *peruviana* *peruviana*

219. *Agrostis* *peruviana*
L. *peruviana* *peruviana* *peruviana*
L. *peruviana* *peruviana* *peruviana*

220. *Agrostis* *peruviana*
L. *peruviana* *peruviana* *peruviana*
L. *peruviana* *peruviana* *peruviana*

221. *Agrostis* *peruviana*
L. *peruviana* *peruviana* *peruviana*
L. *peruviana* *peruviana* *peruviana*

222. *Agrostis* *peruviana*
L. *peruviana* *peruviana* *peruviana*
L. *peruviana* *peruviana* *peruviana*

223. *Agrostis* *peruviana*
L. *peruviana* *peruviana* *peruviana*
L. *peruviana* *peruviana* *peruviana*

224. *Agrostis* *peruviana*
L. *peruviana* *peruviana* *peruviana*
L. *peruviana* *peruviana* *peruviana*

225. *Agrostis* *peruviana*
L. *peruviana* *peruviana* *peruviana*
L. *peruviana* *peruviana* *peruviana*

226. *Agrostis* *peruviana*
L. *peruviana* *peruviana* *peruviana*
L. *peruviana* *peruviana* *peruviana*

227. *Agrostis* *peruviana*
L. *peruviana* *peruviana* *peruviana*
L. *peruviana* *peruviana* *peruviana*

228. *Agrostis* *peruviana*
L. *peruviana* *peruviana* *peruviana*
L. *peruviana* *peruviana* *peruviana*

229. *Agrostis* *peruviana*
L. *peruviana* *peruviana* *peruviana*
L. *peruviana* *peruviana* *peruviana*

fol. lavi. m. ovata.
ad marii Nicaeensis lili & ia Sabuloril.

ss. Lapp

51

52

53

54

55

L. involu
umbell.
Tu

Glauciu
che
in

Cappav

Clema

258. *Euphorbia pinea*
 E. umbell. 5. fid. dichot. involucl. cordat. fol. linear.
 acuminat. confert. caput. breviusculis.
 Agri Nicæensis indigena planta est.
259. ——— *Vegetalis*
 E. umbell. 5. fid. dichotom. involucl. cordat. acut. fol.
 linear-lanceol. super. latioribus.
 in agro Nicæensi.
260. ——— *Serrata*
 E. umbell. 5. fid. 2. fid. dichotom. involucl. diphyll. re-
 niform. fol. amplexicaul. cordat. serratis.
 ubique locis incultis agri Nicæensis.
261. ——— *Spilota*
 E. umbell. 5. fid. 3. fid. 2. fid. involucl. ovat. petal. integr.
 fol. lanceol. subspil. apice serrulatis.
 in Com. Nic. loc. umbrat. & montanis frequentest.
262. ——— *Myrsinites*
 E. umbell. sub 8. fid. 2. fid. involucl. ^{sub} ovatis fol. lanceolatis.
~~mult. spatulatis~~ spatulatis patentibus carnos. mucronat. margine
 scabris.
 Agri Nicæensis indigena planta est.
263. ——— *Dendroidea* V. pl. figur.
 Circa Nicæam in scopulis frequentest.
264. ——— *Characia*
 E. involucl. perfoliat. emarginat. petal. obtus. fol. integerr. hirsut.
 umbell. multifida bifida caule pubescente.
 Tum in convallibus tum locis siccis agri Nicæensis ubiq. copiose.
265. *Glaucium phœniceum*.
 Chelidonium pedunc. 1. flor. fol. sessil. pinnatifid. caul. hirsuto.
 in agro nicæensi inter segetes.
266. *Capparis spinosa*
 Spontanea Nicæa.
267. *Clematis erecta*
 C. fol. pinnat. foliol. ovat. lanceol. integerr. caul. erect. flor.
 5. aut. 4. petalis.
 in collibus agri Nicæensis abunde.

268. *Clematis maritima*
C. fol. pinnat. linearibus cauli simplic. hercagori,
in Sylvis & locis incultis agri Nicæensis.

269. *Fumaria Capnoides*
F. siliquis linearibus 4-gonis cauli diffus. acutangulis,
in Montibus edioribus Comit. Nicæensis.

270. ——— *Spicata*
F. paricarpiis 1.1 permis fol. lobis capillaribus.
Jecur arva Nicæensis.

271. *Polygala monspeliaca*
P. flor. cistitis racemos. cauli erect. fol. lanceolat-linear. acut.
in pascuis siccis nicæensibus
Varietatem P. Vulgaris Credit hanc Cl. Aldrov.

272. *Tritolium subterraneum*
T. capitul. villos. 5-flor. coma central. reflexa rigida
pact. obvolvent.
in agro Nicæensi non varum.

273. ——— *Chevleri*
T. capitul. villos. globos. terminal. solitar. calyc. omn.
fertil. cauli procumb. fol. obcordatis.
Habitat in agro Nicæensi.

274. ——— *Lappaceum*
T. spic. globos. terminal. calyc. set. hisp. rigidi, caule
paulo foliol. ovalibus
in Com. Nic. Jec. prata & agror. marg. & in torrent. tabulor.

275. ——— *angustifolium*
T. spic. villos. conico oblongis dent. calycin. setaceis
subequal. foliol. linear.
Frequentissimum in agro Nicæensi.

276. ——— *stellatum.*
T. spic. pilos. ovat. calyc. patent. cauli diffus. foliol. obcordat.
locis siccis Jecur arva Nicæensis frequenter.

277. ——— *Scabrum*
T. capitul. sessilibus lateral. ovar. Calycibus laciniis inæq.
rigidis recurvis
locis siccis agri Nicæensis.

770 *Tritolium glomeratum.*
Lamiaceae. V. rigid. sessil. calyc. striat. pedicel.

mb.

col.

hem.

269. Clematis maritima
C. fol. pinnat. linearibus cauli simplic. hexagonis
in sulcis & locis. In cultis acribus

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

- 278 *Trifolium glomeratum*.
T. capitul. hemisphaer. rigid. sessil. calyc. striat. patul.
equal.
Locis macilentis & siccis agri Nicaeensis nascitur.
279. — *resupinatum*.
T. spicis subovatis. calyc. inflat. dorso gibbis. caul. prostrat.
in prat. & arv. macilentis, tum siccis usq. Nicaea abunde.
280. — *tomentosum*
T. spic. sessil. globos. tomentos. calyc. inflatis obtusis.
etiam in agro Nicaeensi nascitur haec Trifolii species.
281. — *fragiferum*.
T. spicis subrotundis. calycibus inflatis 2. dentatis reflexis.
caul. repensibus.
in agro Nicaeensi. ad prat. marginis sabulosa torrentium.
282. — *procumbens*
T. spicis ovalibus imbricatis. versic. deflex. persistentibus cauli-
bus procumbentibus.
Nicaeensis agri stirps est.
283. — *filiforme*
T. spic. subimbric. versic. deflex. persistent. calyc. pedicell. caul. procumb.
in prat. Nicaeensibus.
284. *Melilotus italica*.
Trifol. leguminibus racemosis nudis dispersis rugosis obtusis
caule erecto foliol. integris.
In Com. Nic.
285. *Lolus maritimus*
L. legum. solitar. membranac. 4. angul. fol. glabr. bract. lanceol.
in Com. Nic.
286. — *tetragonolobus*
L. legum. solitar. membr. 4. angul. bracteis ovalis
inhabitat agrum Nicaeensem.
287. ~~Lolus~~ *edulis* v. *prol. ligur.*
sec. arva macilentia & ad loca inculta in agro Nic.
288. — *angustissimus*
L. legum. subbinatis linear. strict. erect. caul. erect. peduncul. altern.
In agro Nicaeensi.

289. *Lohet ornithopodioides*
L. legum. subternatis acutatis compressis caulibus diffusis.
Secus arva Nicaeensis abunde.
290. — *Dioicoridis*
L. caul. erect. ramor. pedunc. subflor. legum. teret. ovalis forat.
ad maris ripas in arenosis herbidi. prope Hospitium sic dictum
Villa paucis repertus est.
291. — *hirtulus*
L. capitul. subcylindric. caul. erecto hirsuto legum. ovalis
horis incultis & in sylvis apparet Nicaeas. vulgaris.
292. — *recta*
L. capitul. subglobos. caul. erect. laevi legum. rect. glabris.
Nicaea ad sepes & pratorum margines.
293. — *Cyathoides*
L. capitul. dimidiatis caule diffuso ramorissimo fol. tomentosis.
in arid. Nicaeensibus mari proximis.
294. *Dorychnium monspeliense*
L. capit. asphyllis fol. sessil. 5 ual.
abunde in Com. Nic.
295. *Anthyllis tetraphylla*
A. herbacea fol. quaternopinnat. flor. lateral.
Secus arva macilenta Nicaeensis.
296. — *barbajovis*
A. pilicosa fol. pinnatis aequalibus tomentosis flor. capitatis
ad abruptis rupes mari Nicaensi proximae.
297. *Buceva polyceraton. Trigonella?*
Folia terrena versat. coroll. alit subaequale cavina brevissima
fructus est siliqua compressa falcata.
J. legum. subcylindric. congest. erect. suberect. longilincar. pe-
dunc. mult.
in agro Nicaensi.
298. — *corniculatum*
J. legum. pedunc. congestis declinat. subfalcat. pedunculo long.
subspinoso caule erecto
in Agro Nicaensi.

2. fus

2. h. tenu
2. h. tenu

2. h. tenu
2. h. tenu

2. h. tenu

2. h. tenu

2. h. tenu

2. h. tenu

2. h. tenu

2. h. tenu

270. *Leptocarpus* ...
... ..
... ..

271. *Leptocarpus* ...
... ..
... ..

272. *Leptocarpus* ...
... ..
... ..

273. *Leptocarpus* ...
... ..
... ..

274. *Leptocarpus* ...
... ..
... ..

275. *Leptocarpus* ...
... ..
... ..

276. *Leptocarpus* ...
... ..
... ..

277. *Leptocarpus* ...
... ..
... ..

278. *Leptocarpus* ...
... ..
... ..

279. *Leptocarpus* ...
... ..
... ..

280. *Leptocarpus* ...
... ..
... ..

281. *Leptocarpus* ...
... ..
... ..

282. *Leptocarpus* ...
... ..
... ..

Bucera
Maring
Ch
Co
Ch
H
Ch
Co
8

299. *Bucera monipeliacum*

J. leg. sessil. congest. arcuatis divaricatis inclinatis bre-
vibus peduncul. mucronato inermi.
in sterilibus agris Nicaensibus.

300. — *fornum graecum.*

J. legum. sessil. strict. erectiscent. subsalsatis acuminatis caule
basivamolo stolonifero
ad agrorum margines in Com. Nic.

301. *Medicago radiata*

Ch. leguminibus venispermis marginedentatis fol. ternatis.
in agris Nicaensibus.

302. — *orbicularis*

Ch. legum. solitar. cochleatis depressis planis stipul. ciliatis caule diffus.
Copiose sedes agris in Com. Nicaensibus.

303. — *scutellata.*

Ch. cochleata major dicarpos capsula rotunda globosa scutellata
Frequent in Com. Nicaensibus inter seget. & secus arva.

304. — *intertexta.*

Ch. cochl. major. spinosa dicarpos capsulae spinis longioribus sursum
& deorsum tendentibus
Haec etiam Nicaensibus agris indigena est.

305. — *arabica*

Ch. cochl. minor polycarpos annua capsula majore alba folio
cordato macula fusca.
In agro Nicaensibus inter segetes.

306. — *lubbicata*

Ch. legum. cochl. inerm. subcylindr. erectum. complanatis.
inter segetes in agro Nicaensibus.

307. — *ciliaris*

Ch. legumin. congest. globos. hispida. stipul. ciliatis.
Cum praecedentibus in eodem locis.

308. — *vigridula*

Ch. biphyll. legum. cochl. spinosa foliol. inferior. cuneiform. velut.
superioribus subrotundis.
In agro Nicaensibus.

309. — *muricata*

Ch. cochl. dicarpos capsula spinosa rotunda minore
etiam in agro Nic. nascitur.

310. *Medicago laciniosa*
M. legum. coroll. spinos. stipul. contort. fol. oblong. duplicat. serratis.
 In agro Nicaeensi.
311. ——— *marina. V. pl. figur.*
 in sterilibus litor. maris proscimit.
312. *Ononis antiquorum.*
O. flor. solitar. foliol. major. fol. inferior. ternat. v. var. laeviuscul.
spinol.
 in agro Nicaeensi.
313. ——— *Cherlevi*
O. pedunc. 1. flor. avistat. fol. ternat. stipul. serratis.
 In Jaspuloni & ad rupes Villae francae Nicaeensi.
314. ——— *Vitcola*
O. pedunc. 1. flor. avistat. fol. simplic. infim. ternat.
 locis aridis circa Menton et inter Roche Tallinde & Biot.
315. ——— *pinguis*
O. pedunc. 1. flor. avistat. fol. ternat. lanceolat. stipul. integerr.
 in collibus Nicaeensibus.
316. *Cytisus lupinus V. pl. Tend.*
 inter Jaspello & Lanteca.
317. ——— *gracilis*
C. fol. simplic. lanceolat. lineav. v. var. angulatis.
 Nicae exemplar lectum habet Cl. Allione.
318. *Genista sagittalis*
G. v. var. angustifolia membranac. articulatis fol. ovato-lanceolatis.
 inter Jaspello & Lanteca.
319. ——— *hispanica*
G. spin. decompositis v. var. flor. fer. inerm. fol. lineav. pilos.
 in aridis apricis Com. etc.
 Dubitat. Cl. Allione num vere a Germanica differat.
320. ——— *candicans*
G. fol. ternat. subt. villos. peduncul. lateral. sub. 5. flor. foliol. legumin. hirsutis.
 in montibus Nicaeensibus Calidioribus.
321. *Hedysarum spinosissimum V. pl. figur.*
 in agro Nicaeensi loc. Jaspello maris prosc. & securaria.

322. Onobrychis sarcocolla V. pl. ligur.
in locis aridis et silicibus con. nicaens.

310. *Medicago laciniosa*

M. lequum var. *laciniosa*. Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3

311. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3
312. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3
313. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3
314. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3
315. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3
316. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3
317. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3
318. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3
319. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3
320. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3
321. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3
322. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3
323. *Medicago laciniosa*
Rhizom. contort. fol. oblong. duplicat. 1/2 v. 1/3

322. *Onobrychis sarcocolla* V. pl. figur.
inter montes apud Con. nicænsis.

323. — caput galli
Medicago. fol. pinnatis legum. monosperm. cristat. aculeat.
pet. equal. caul. diffus.
ad Colles aridos & sabulosos torrentium in agro Nicænsi.

324. *Vicia* ~~Nica~~ *Bythynica*
V. legum. pedunc. solit. erect. foliol. quaternis ovali-lanceol.
hipoc. dentatis.
Nicænsis agri indigena planta est.

325. — *Lathyrus*
V. Leg. sessil. solitar. erect. glabr. foliol. den. inferior. obcor.
dat.
in agro Nicænsi.

326. — *peregrina*
V. leg. subsessil. pendul. glabr. 4. sperm. foliol. linear. emarginat.
inter seget. agri Nicænsis.

327. *Ervum* *monanthos*
L. pedunc. uniflor.
agri Nic. indigena planta est.

328. — *erilia*
L. germin. undat. plicat. fol. impari. pinnat.
inter seget. Agri Nic. occurrat.

329. *Lathyrus* *licera*
L. pedunc. uniflor. circ. diphyll. legum. ovat. compress.
obso. canaliculatis
in agro Nicænsi. spontan.

330. — *laticul*
L. pedunc. 1. flor. circ. diphyll. 4. phyllis. legum. ovat. compress.
obso. bimariginatis.
etiam Nic. agri indigenus.

331. — *setifolius*
L. pedunc. 1. flor. circ. 2. phyll. foliol. setac-linear.

332. — *angulatus*
L. ped. 1. flor. aristat. circ. diphyll. simpliciss. foliol. linear.
Vulgaris in agro Nicænsi.

333. — *heterophyllus*
L. pedunc. multiflor. circ. diphyll. 4. phyllis. foliol. lanceolat. internod.
membranaceis.
in umbratis di lucevane.

334. *Pisum maritimum*
 sp. petiolis supra planiusculis cant. angulatis. stipulis sagittatis pedunculatis floribus.
 In agro Nicaeensi nascitur.
335. ——— *ochrus*
 sp. petiolis decurrentibus membranaceis. diphyllis pedunc. 1. floribus
 inter segetes Nicaeenses.
336. *Coronilla juncea*
C. fruticosa fol. pinnatis ternatisq. linear-lanceolatis subcarnosis
 obtus.
 in agro Nicaeensi
337. ——— *Valentina*
C. fruticosa fol. subovatis stipulis suborbiculatis.
 in colle di Montalbano.
338. ——— *glauca*
C. fruticosa foliis suborbiculatis stipulis ovatis pedunc. complanatis
 in agro Nicaeensi ad super.
339. ——— *securidata*
C. herbacea legum. falcato-glabratis foliis plurimis.
 in agro Nicaeensi vulgariter est.
340. *Ornithopodium compressum*.
O. fol. pinnatis legum. recurvatis compressis rugosis bracteis pinnatis
 in agro Nicaeensi.
341. ——— *scorpioides* N. sp. figur.
Securaria maritima agri Nicaeensis.
342. *Scorpiurus vermiculata*
 Gen. Calice. semi-5-fido dentibus unguibus minutis divitis Vaseillum
 reflexum subrotundum emarginatum patens alae oblongae laevae Carina
 recta gibba ungue divisa Siliq. teres in se contracta articulat. rugosa vel hispida
 J. pedunculatis 1-floris legum. fectis & indigne squam. obtusis.
 inter segetes & ad vias in agro Nicaeensi.
343. ——— *muricata*
J. pedunc. 2-flor. legum. erectissimum obtuse aculeatis
 similibus locis prius nascitur.
344. ——— *subvillata*
J. pedunc. sub 4-flor legum. erectiss. spic. confectis erectis.
 Nicaeentem agrum incolit.

Spital

ne. 1/2

at. 1/2

compleat

mit.

black. 1/2

Vanillan
laste. 1/2
1911. vel. 1/2
1911.

1/2

1/2

[Faint, illegible handwriting throughout the page, likely bleed-through from the reverse side.]

1. The first part of the paper is devoted to a general
discussion of the subject, and to a statement of the
object of the investigation.

2. The second part is devoted to a description of the
materials used, and to a statement of the methods
employed in the investigation.

3. The third part is devoted to a description of the
results obtained, and to a statement of the conclusions
drawn from them.

4. The fourth part is devoted to a description of the
apparatus used, and to a statement of the results
obtained from it.

5. The fifth part is devoted to a description of the
conclusions drawn from the results obtained, and to a
statement of the conclusions drawn from them.

345 Hipp
Gen
intra
intra
intra

346 —

347 Spod

348 —

349 / Ma
Gen
intra
intra
Trib
in C

350 Cede

351 Ana
Gen. P
Ala
Furin
Ci
Joffe
tagor
nide
Vulva

352 Alva
Gen
obv

345. *Hippocrepis unisiliquosa*
 Gen. Essentialis. Character Constitit in siliqua compressa longa incurva tubera
 interiore ad medium usq. inter sem. sinuata sola & equina ad instar se-
 mina incurva.
 H. legum. sessil. solitar. vact.
 Abunde in Com. Nicaeni.

346. ——— *multisiliquosa*
 H. legum. pedunculat. confert. ^{circularibus} ~~areolatis~~ marg. ^{altero lobatis} ~~retusis~~ ^{reflexis}.
 etiam haec sed rarius in agro Nic. occurrat.

347. *Spartium purgans*
 J. ram. teret. striat. fol. lanceol. sessil. pubesc.
 freq. et. in mont. calidior. Com. Nicaeni.

348. ——— *spinosum*
 J. fol. ternat. ram. angulat. spinos.
 in Com. Nic. uberrime crevit.

349. *Psoralea bituminosa*
 Gen. Calyce. 1. phyll. 5. fid. irregul. tubercul. asperus. Vexill. subrotund.
 emarginat. altiuscul. ala oblonga parva. Carina dipetal. lunulata siliqua
 brev. calyce contenta semen 1. aut 2. contin.
Trifolium bitumen vedolens
 in Comit. Nicaeni ubique.

350. *Cercis siliquastrum*
 C. fol. cord. orbiculat. glabri.
 frequent in sylvis et ad rupes in agro Nicaeni.

351. *Anagyris foetida*.
 Gen. Calyce. campanul. 5. dentat. irregul. Vexill. breviss. rectum.
 Ala oblonga vexillo longiore. Carina 2. petal. longissima Stamina
 omnia distincta siliqua teretiuscula magna semina reniformia.
 Circa Nicaenam ad rupes
 suffrutesc. fol. altern. petiol. ternata foliol. sessil. oblong. acuminat.
 haecc. supra virid. subt. albicans. odor. grav. & fetid. pedunc.
 arbor. multifl. binis-ternis-4. foliol. oblongis Cal. 5. fid. glauc.
 & velut argentea. Vexill. aureum coctus. petala pallidiora.

352. *Alvagalus hamolus*.
 A. caulescens procumbens legum. tubulat. recurvat. glabris foliol.
 obcordat. subtus villosis.
 Locus Larcois Collium Nicaenium.

353. *Bilevula pelecini* L.
 Gen. Calyce monophyllas semib. fid. irregulari. Vexillum majus late-
 ribus reflexis alae oblique vexillo breviores Cavina obliqua siligina com-
 pressa 2-loculari septo valvi contrario semina reniformia.
 Nicotia inter segetes non rara.
 Caul. herbac. ramos. fol. altern. imparipinn. foliol. mult. opposita
 subseriat. obcordat. integerr. flor. axill. solitarii peduncul. dimidio
 capituli rubelli.

354. *Eragium maritimum* V. pl. ligur.
 in sabulo ad mare Lintus

355. *Echinophora spinosa* L.
 Gen. involucri pauce folior. involucri. 1-phylli. multif. flos. radiat.
 pet. inaequal. sem. sterilia coronata in involucrio indurata.
 E. foliol. subulato-spinosis integerrimis.
 frequent ad Lintus Maris Nicotensis.
 Caul. er. b. t. ramos. striatus subulato. fol. altern. 2-pinnatif. subulato
 lacin. linear. apice spinos. pedunc. later. subulato. pet. alb. sub. pur-
 purascentia.

356. *Psallidula oppositifolia* V. pl. ligur.
 in agro Nicotensi locis umbratis & montibus asperis.

357. *Fevula Communis*. Gen. Semen. magn. oblong-ovate 3-striat. cinch.
 ala magna solida non membranacea.
 F. foliol. linear. longior. simplicibus.
 In agro Nicotensi nascitur.
 Albidus humana radice crassa lactescens fol. alterna. amplexi-
 2-pinnat. umbell. robust. pitat. prae. vestesca lutea.

358. — *nodiflora* V. pl. ligur.
 in Monte Joet Comit. Nicotensis.

359. *Pencedanum minus* V. pl. Tend.
 inter la Ghiaudola & la Scavena.

360. *Selinum Carvifolia* Chabraz
 J. fol. multiplic. radical. circa costam decussati
 in pratis Nicotensibus

361. *Thapsia villosa* Gen. Flor. equal. petal. integri lanceolati; invo-
lucrum nullum. Sem. oblong. 3-lobatum. magna ala circumscripta lu-

353. *Bilevula pelecini* L.

Com. Callus monophyllus semib. p. d. irregulari. Versillum majus late.

35

3

35

35

35

35

36

361. *Thapsia villosa* Gen. Flor. opposit. petal. integris lanceolatis involu-
crum nullum. Sem. oblong. 3-lobatum magna ala cinctum quae su-
pra & infra ampliter truncata est & emarginata.

T. foliol. dentat. Villob. bar. (caudiculis)
in agro Nicotenti.

Caul. ca. 6 ft. & ultra brachiatus fol. crassiusculis & aspera in ala
oppositis subjecta segment. longior. semunitat. umb. lat. pet. futa.

362. *Lactepitium Gallicum* V. pl. Vall. Mur.
inter Tenda & Eleavena.

363. *Ligusticum speloponeriacum.*

L. fol. multiplicato pinnatis foliol. pinnatis incisis.

in com. Nicotenti. circa Toppello & La Ghiaudola.

invol. Univ. 5-6. phyll. parial. 2. aut 3. phyll. petal. cordat. pubescent.
Sem. glab. oblong. non rostrat. profund. sulcat. live 1-lobat.

364. *Coriandrum testiculatum.* V. pl. lignv.
in agr. com. Nicotenti.

365. *Pimpinella peregrina*

P. fol. radical. pinnatis crenatis sum. cuneiform. incisis.

L. Beckard: in sterilibus Nicotenti. bus ad prothum d. pimpin.

366. *Seseli annuum*

S. petiol. rament. membranac. ventricol. emarginat.

in collibus com. Nicotenti. frequens est.

367.

glaucum

S. petiol. rament. membranac. oblong. integris foliol.
singulis binatisq. cuneatis. latius petiol. longior.
in collibus aridis agr. Nicotenti.

368.

tormentosum

S. caul. alto rigido foliol. linear. fasciculat.
inter Lucerne & La Scarenna ad vixet.

Se Seligenus flores hab. florulor. pedicel. involucri. parial. poly-
phyll. universale vix ullum aut 0. petala cordata reflexa fructus
subrotundus aut ovatus. Sem. ovat. acuminat. intus planungitum
h. corbatur.

369.

Oenanthe globulosa Vid. Gen. ad pl. lignv.

O. fructibus globosis

in herb. d. & sub hum. agr. Nicotenti.

370.

pimpinelloides V. pl. lignv.

part. loc. sub hum. d. ex graminor. in com. Nicotenti.

371. *Supplementum volundifol.*
in agro Nicotianum collinum.
372. — *Jemiconoposium*
fl. caule herbaceo fol. lanceol. umbell. acillar. terminalibus.
sem. scabris
Circa Villa Paucina in sterilibus vasis Saint Office.
373. — *juncum v. pl. ligur.*
Habitat in agro & Com. Nicotianum.
374. — *temuissimum*
fl. fol. angustissim. umbell. simp. altern. 5-phyl. sub 3-fl.
in agro Nicotianum.
375. *Carum Bunium*
Gen. involucri. utriusq. dext. petal. cord. aeq. incis. sem. oval. profund.
striata intus plana.
Cum bell. & fid. fol. radical. bipinnat. linear. caul. sub erectis
in Vineis Savorgii & Circa Lucerne.
376. *Sison anisum.* Involucri. pinnatif. sub 4-phyl. petal. collat.
aequal. n. oval. oblong. parva sem. sub biliter striata.
Carum Bunium & Sison ad finem sunt gen. ut digne disting. possint.
fl. fol. pinnat. umbell. erect.
in Com. Nicotianum.
377. *Cistidium maritimum v. pl. ligur.*
in mari Vetusis. Com. Nicotianum.
378. *Sium falcaria.*
Gen. fl. aeq. fessit. petal. aequal. cordat. fruct. oblong-oval.
sem. oval. seuster striata intus plana. calyc. minim. coronata.
fl. foliolinear. decurrent. connatis.
in agro Nicotianum.
379. — *nodiflorum*
fl. fol. pinnat. umbell. acillar. sessil.
Cl. Bellard: prope Sappello ad muros Sublimisidos.
380. *Myrrhis australis*
fl. sem. tubulatis bipid. flor. radicalibus caul. laxis.
in agro collinum Nicotianum.
381. *Daucus mauritanicus*
fl. sem. bipid. florum. cent. ster. carnol. receptacul.
comm. hemisphaerico
in pratis agro Nicotianum.

382. *Dorcas bipartitus*.
D. bipartitus central part: nullo caule lucido.

1891

2. $\frac{1}{2}$ of the total

Dep. 2401. / 3/10/11

cond. No. 1221

Mytilus parasiticus

ix di King - 1/12

Blang-ai
mim-ai to fa.

1000/1000.

625.64

st. verapn cul.

371. *Supplementum Schmidt. fol.*
in aq. Nicotianum Collium.

382. *Donneri hispanicus*.
D. sem. aculeatis centrat. ster. i. nullo caule lucido.
in agro Nicotensi.

383. *Caucalis latifolia*
Involuc. etc. foliol. non pinnatifid. sed integ. v. paucifol. quod
interdum deest sem. striat. aut costat. fect. aculeis aut setis rigidis
uncinatis.
C. umbel. unives. 3-fida partial. 5-spermi. fol. pinn. serratis.
in agro collium Nicotensium.

384. — *maritima*
C. umbel. 2-fid. umbellat. 2-spermi. involuc. semine
brevior. linear.
in agro Nicotensi.

385. — *nodiflora*
Tardylum umbell. simplic. Jectil. sem. exterior. hisp. d. i.
ad limit. agro. & in partibus latioribus C. Nicotens. frequens.

386. — *nodosa*. V. pl. lign.

in vineis & ster. limitibus Nicotens.

387. *Geranium ciconium*
G. pedunc. multiflor. 5-andr. fol. pinnat. pinnatif.
obtus.
Nicot. ad semitas.

388. — *moschatum*.
G. ped. multif. flor. pentandr. fol. pinnat. incis. cotyledon.
pinnatifidit.
Nicot. vespertur

389. — *malacoides*
G. ped. multiflor. flor. pent. fol. cordat. sublobat.
Circa Nicotam ovitur.

390. — *maritimum*
G. pedunc. sub 3. flor. flor. pentandr. fol. cordat. incis.
crenatis scabris caul. procumbentibus.
Junct. vias Nicot. ad loca maris proxima.

391. — *lucidum*
G. ped. 2-flor. calyc. pyramidal. angulat. elevato rugosis
fol. 5-lobis robustis
in agro Nicotensi.

392. *Malva parviflora*

Calyce exterior 3phyllus interior semi 5. fidus.

Herb. caule patulo fol. angulat. flor. axillar. sessil. glomeratis. Calyc. glabri patentibus.
in Comit. Nicæensi.

393. ——— *Nicaeensis.*

Herb. caul. pilos. decumbent. herbac. fol. subcordat. 5. lobis acutis.
pedunc. erectis.
Cl. Bellardi in agro Nicæensi prope portum di. Rimpia.

394. *Lavatera arborea.*

Differt. Genus hoc. a Malva calyc. exterior. 3. fido non 3phyllus.

Herb. caule arboreo fol. 7. angulat. tomentos. plicatis.
pedunc. confert. 1. flor. axillaribus.

Reperitur ad super. liton. Nicæensi.

395. ——— *olbia*

Herb. caul. pilos. fol. 5. lob. dentat. pedunc. solitari.
Nicæensem agrum inhabitat.

396. ——— *thuringiaca*

Herb. caule herbaceo fruct. denudatis calyc. incisis.
in Comit. Nicæensi.

397. ——— *trimestris*

Herb. fol. glabris caule scabro herbaceo pedunc. 1. flor.
fruct. orbicul. sect.

in oliveis Villafranca Nicæensi reperitur.

398. ——— *maxima.*

Herb. caule pilosissimo lanato fol. cordat. orbiculat. 5. lob. obtus. crenat.
pedunc. 1. flor. stipulis sericeis deciduis.
ad super. liton. & agri Nicæensi abunde.

399. *Alcea Rosea.* Calycem exterior. habet. semi 6. fidum.

Al. fol. sinuatis angulosis
in Comitum Nicæensi.

an Althea hirsuta Cal. erecta. plusquam 5. fid.
lucida & ibi c. crenat. ceter. 3. fid.

392. *Malva parviflora*
C. h. bidus.

39

39

39

396

397

398

399

400. *Althaea hirsuta* Cal. erect. plurigam. 5. fid.
A. fol. inter. tuberos. ib. 70. cruent. ceter. 3. fid.
inhabit. super & Collet Com. Nicaeni.

401. *Hypericum tomentosum*.
H. fl. trigyn. calyc. serratis glandul. fol. semi am.
pleuricant. flexuos. tomentos. caul. prostrat.
ad fustes rivulos & aqueductus in agro Nicaeni.

402. *Croton tinctorium*. Monoezia monadelphica.
Gen. Masc. calyc. cylindr. 5. dentat. coroll. 5. petal. Stam.
10-15.
Fem. Cal. polyphyll. Corol. 0. Styl. 3. bifid. Capsul. 3. locul.
sem. 1.
C. fol. rhomb. repandis caput. pendul. caul. herbac.
in Agro Nicaeni: frequens.
Planta tenella pulverulenta caul. cubital. variet. fol. altern.
long. petiol. sinuata nervosa fl. axillar. congesti pedunculat.
calyc. 5. phyll. petal. lanceol.

403. *Ranunculus muricatus*
R. sem. aculeatis fol. simpl. lobat. obtus glabr. caul. diffus.
in agro Nicaeni.

404. ——— *parviflorus*. V. fl. ligur.
in arvis subhumid. Com. Nicaeni.

405. *Corrigiola litoralis* V. fl. lig.
locis arenosis mar. proximis.

406. *Delphinium peregrinum*
D. nectar. diphyll. flor. solitar. fol. multipartit. foliol.
linear. acuminatis.
Circa Nicaenam nascitur.

407. ——— *Staphysagria*
D. nectar. 4. phyll. petal. brevior. fol. palmat. lob. obtus.
Reperitur Nicaea.

408. *Nigella damascena*
Agri Nicaeni: familiaris planta.

409. *Helioscopus niger*.
H. scapo subbifloro subundo fol. pedat. / hore vireo
in frigidioribus Nicæa montibus.

410. *Saxifraga granulata*
J. fol. caul. n. reniform. lobat. caul. ramor. radic. granulata.
Loci? in p. h. h. h. Com. Nicæa.

411. *Saxifraga vaccaria*
J. calyc. pyramidalis 5. angulat. fol. ovat. acumin. sessil.
in Com. Nicæa. inter leges.

412. *Velexia rigida*. Calyc. filiform. cylindr. 5. dent. Corolla parva
5. petala / tam. 5. vel 6. styl. 2. Caput. 1. locut. cylindr. polyperm.
Stam. in simplic. serde.
in h. v. h. h. ad limitey agerum Nicæa.

Caul. dequivalvar. Vannolitt. duri. fol. connat. tubulat. integerr.
pilosa. fl. axillar. solitar. sessile. Corol. limbus minimus.

413. *Silene anglica*
J. h. m. petal. emarginat. fl. erect. fruct. reflex. peduncul.
altitud.
Secus arva Nicæa.

414. ——— *ceratoides*
J. h. v. pet. emarg. fruct. h. c. erect. calyc. subpilol. sessil.
Medic. Cauvini in agro Nicæa. legib.

415. ——— *S. Vulnera*
J. petal. integerr. subrotund. fruct. erect. altern.
Secus arva Nicæa.

416. ——— *nocturna*
J. fl. spicat. altern. secund. sessil. pet. bifid.
in agro Nicæa. Oliveb. passim occurrat.

417. ——— *conoides*.
J. calyc. fruct. globos. acuminat. styl. 30. fol. glab. petal. integ.
Nativum in agro Nicæa.

418. ——— *noctiflora*
J. calyc. 10-angul. dent. tub. æquant. caul. dichot. petal. bifid.
in Com. Nicæa.

419. ——— *Nicæensis*
J. v. h. v. fol. linear. obtus. petal. renib. fruct. ovat. calyc. 10-
in agro Nicæa. Secus le vas frequenter.

116

annulata.

117. 118.

ab. Corolla
cylindric. po

ab. lat. int
mispum.

refl. pet

119.

ab. petal. 11

ab. petal. 11

ab. oval. cal. 11

petal. 11

400. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 401. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 402. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 403. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 404. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 405. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 406. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 407. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 408. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 409. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 410. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...

400. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 401. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 402. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 403. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 404. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 405. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 406. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 407. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 408. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 409. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...
 410. *Valeriana anglica*. Calyx. Rotundum. Stylus. 5 denis. Fructus. ...

420. *Silene inappetita*.
S. caul. dichotom. panicul. calyc. laev. petal. breviss. emargin. fol.
glabra lanceol.
Hoc etiam Nicæensis agri indigena est.
421. *Agrostemma ciliolata*
A. glabra fol. linear. lanceol. petal. emargin. coronat.
in Comitatu Nicæensi.
422. *Tamariacus gallicus*
T. flor. pentand. fol. viridib. tenuibus
in agro Nicæensi.
423. *Statice limonium*
S. scapo paniculato tereti fol. laev. enerv. subt. mucronat.
agrum Nicæensem ad maris litus inhabitat.
424. — *cordata*
S. scapo panicul. fol. spatul. velut.
ad vases maritimos frequent.
425. *Frankenia pulverulenta*.
calyc. subcylindricus 5-dent. persistent. lثم. 6. petal. 5. styl. 1.
hygmat. bifido caput. 1. locut. ab. im. adum. in 3. valvul. dehisc.
Polysperma.
f. f. obovat. velut. subt. pulverat.
in agri Nicæensi arenos.
caul. humifus. palmat. ramor. rubent. fol. oppos. vel. quatern. fascic. erigui
amæthistini.
426. *Acerda alba*
A. fol. pinnat. fol. ovibus tetragynis calyc. seropassibil.
in C. Nicæensi. locis sterilibus & arenosis.
427. *Tribulus terrestris* V. pl. ligur.
Nicæa securaria abunde.
428. *Halmala multifida*. *Pegannum L.*
cal. 5. phyll. coroll. 5. petal. aequal. lثم. 15. styl. 3. fid. caput. 3. locut.
Polysperma.
f. fol. multifidi.
in agro Nicæensi nascitur.
caul. cubital. ramor. /wizhu. fol. alterna terrib. succulenta Flor.
oppositifol. solitar. pedunculati fol. Calycin. linear. 3. longior.
petal. luteo virid.

429. **Cistus**. Monipeliensis.

C. arboreus, erectus, fol. lineari-lanceol. sessil. utrinq. villis.
3-nerv.
Freg. in arid. incult. et Jaceo. circa Villanueva Coppori.
circa Berra in Com. Nicoteni.

430

Salvifolius.
C. arbor. erectus, fol. ovato. petiolat. utrinq. hirsutis.
in agro Nicoteni ad super aridas.

431

incanus.
C. arb. erectus, fol. spatul. toment. rugos. inferior. basi vaginatis. connatis.
in Com. Nicoteni. non in freg.

432

medius
C. fraxinea, folio Salviae flore ochrea colore
in agro Nicoteni.
Caul. rubent. viscos. pedunc. solitari. axillar. & terminal.
fol. petiol. multum rugos. virid. ovato-lanceol. in amb. dentatis.

433

albidus V. pl. liquor.
in agro Nicoteni ad super aridas.

434

crispus
C. arbor. erectus, fol. lanceol. pubesc. 3-nerv. undul.
in Comit. Nicoteni.

435

laevipes.
C. suffrutescens, adscendens, erectus, fol. altern. fascicul.
filiform. glabr. pedunc. racemos.
in agro Nicoteni frequent.

436

canus
C. suffrutescens, erectus, procumbens, fol. opposit. obovat.
villat. subt. toment. fl. subumbellat.
Habitat in C. Nicoteni.

437

italicus.
C. suffrutescens, erectus, fol. opposit. hirsut. inferior. ovato. superior.
lanceolat. v. n. parent.
in rupibus circa Lucerame.

38 *Cistus tuberosus*
 C. eschschol. perenn. fol. radic. ovab. 3. nerv. toment. caulin.
 alab. lanceol. humil. alternis.

alt. 1.5 m.
 in ...

in ...

in ...

terminal.
 in ...

alt.

terminal.

alt. ovab.

alt. superior.

alt.

alt.

alt.

alt.

429. *Cistus monspeliensis*.

C. arboreus ex Hippocr. fol. linear-lanceol. sessil. utrinq. villis.

43

43

43

43

434

435

436

437

438. *Cistus tuberosus*
C. erectus. stipul. prostratus. fol. radice. ovato. 3. nerv. tomentoso. caulibus
glabris. lanceolatis. summis alternis.
in Com. Nicaeensi. Sterilibus & ruppibus frequenter
circa Nicaeam.

439. ——— *guttatus*. V. M. Liguor.
in paludibus siccis Sterilibus Nicaeensi. Comit.

440. ——— *leucifolius*
C. herbaceus. glaber stipulatus. flor. solitarii suberecti. fol. ternati.
oppositi.
in Com. Nicaeensi.

441. ——— *laticifolius*
C. herbaceus. patulus. villosus. stipulatus. flor. racemosi erecti.
pedunc. horizont.
in Com. Nicaeensi.

442. ——— *thymifolius*
C. suffrutescens. stipul. prostratus. fol. ovato-linear. oppos. breviss.
congest.
in collibus maritimis frequenter.

443. ——— *roseus*
in Com. Nicae.

444. ——— *romarinifolius*
C. hirsutus suffrutescens stipulatus. fol. ovato. calyc. hirsutus. lin.
in ar. di. Circa Villafraancam Nicaeensi.

445. ——— *glutinosis*
C. suffrutescens. stipulatus. fol. linear. oppos. alternisque
pedunculatis. villosis. glutinosi.
in rupibus agri Nicaeensis non infrequenter.

446. *Gavideella nigellae* Wurm. Decandria 3-gynia
Cal. 5. phyll. coloratus petala 5. aequalia. 2. lab. nectariifera caps. 3. connatis. polysper.
G. fol. tenuiss. divisi.
Nascitur in agris Nicaeensibus.

447. *Telephium implexati*
 Calyce 5-phyll. petal. 5. equal. Stam. 5. Styl. 3. fr. 3. Capitul. 1. locul. 3-valv.
 Polysp. J. fol. alterni/
 frequent in Comit. Nicæensi
 Caulis procumbens teret. foliis fol. obverse ovata glauca nervis in caulem
 decurrentibus ima robustiora fl. terminal. umbell. alb.

448. *Linum perenne*
 l. calycibus capsul. 2g. obtus. fol. altern. lanceol. integerr.
 in montibus circa Sospello & Lucerame.

449. ——— *hirsutum*. V. pl. lig.
 in Com. nicæensi. spatium occurrat.

450. ——— *marbonense*.
 l. calyc. acuminat. fol. lanceolat. sparsi. Hict. scabr. acuminat.
 Caul. teret. basi ramos.
 frequent in sarcois inter Lucerame & Touet C. Bellardi in mon.
 tib. a Castiglione ad Menton & a l'Escavene ad Breglio.

451. ——— *hirsutum*
 l. calyc. tubul. fol. lanceolat. Hict. mucronat marg. scabris.
 locis arid. in agro Nicæensi uberrime.

452. ——— *campanulatum*
 l. fol. bas. utrinque punctat. glandulosa.
 inter l'Escavene & Drapp.

453. ——— *flavum*.
 l. calyc. suberrato scabris lanceol. suberectilib. panicul. ram. dichot.
 in collibus maritimis Nicæensibus.

454. ——— *maritimum*
 l. calyc. ovat. acut. n. utricul. fol. lanceolat. inferior. opposit.
 in humidis & stagnantibus agr. nicæensis & præcipue l'Escav.
 fl. le Var.

455. *Polycarpon tetraphyllum*
 Cal. 5-phyll. petal. 5. persisten. cal. minor. Stam. 3. Styl. 3. Capr. uniloc. 3-valv.
 Nicæensis agr. locis cultis / monte natali
 Caulis semipalmat. glaber prostratus ramosus fol. elliptica obtusa
 integerr. bina in caule a 2 ramov. originem quaternam flosc. albi.

Hand. 30

456 *Arenaria marina*

A. l. lina - linn. l. lina - l. lina - l. lina

l. lina

l. lina

l. lina

l. lina

l. lina

l. lina

l. lina

l. lina

l. lina

l. lina

447. *Telephium implexati*
Calyc. 5. coroll. 4. equal. Stam. 5. Styl. 3. pist. Capit. 1. lobul. 3. unal.

44

45

46

47

48

49

50

51

11629 2. *Arenaria marina*

A. marina - *Islandia* - *Reykjavik* - *Reykjavik*

447. *Telephium imperati*

Stem 6. Root 2. Lvs. Capit. 1. Lvs. 3. Lvs. 4.

4.

4.

4.

4.

4.

4.

4.

4.

456. *Arenaria marina*
A. fol. linear. longitudine internodiis
prope portum Villafrancae uicinis.

457. *Sedum Nicaeense*.
S. fol. subulatis, parvis basi solutis, alternis cyma bifida flor. long. pedunculatis
Rarissima planta nascitur in scopulosis prope Savorgio.

458. *Coriaria myrtilifolia*. Dioecia decandra
Male. Cal. 5. phyll. Corol. 5. petal. calyc. limbitum connexa anther. 2. styl.
Fem. Cal. 5. Corol. maius styl. 5. sem. 5. petal. succulent. baccalis tecta.
C. fol. ovata oblongis
ad sepes in Com. Nicaeensi Vulgaris planta est.
Caul. pubescens 2. cubit. ligni cort. cinereo maculatis fol. pinnatis
foliol. 3-nerv. bractae ovatae racemiformi Cor. parva Vivida
fem. gaud. anth. steril.

459. *Rhamnus alaternus*
R. inerm. fl. dioic. stigmat. 3. foliis fol. serratis.
in agro & collibus Nicaeensi abunde.

460. ——— *R. alaternus*
R. aculeis geminis inaequal. fol. binervi. subternat. peduncul.
solitar. fruct. marg. membranaceo clypeiformibus.
incolit Comit. Nicaensem.

461. ——— *Zizyphus*
R. aculeis geminatis altero recurva flor. digyn. fol. ovata oblong.
glabris.
Habitat in Comit. Nicaensi.

462. *Melia Azedarach*
Calyc. 5. dent. petiol. in med. flor. nectar. cylindric. multidentat.
10. anther. latine breviss. pedicelatis styl. 1. fruct. drupae firma.
Circa Nicaeam collectam habuit Cl. Allouin.
Fol. parcinervis. altern. petiol. foliol. petiol. integr. fl. axillar. ra-
cemosi longe peduncul. parvi purpurei.

463. *Prunus Mahaleb*.
P. flor. terminal. corymb. fol. ovata.
in montibus Nicaensi Comitatus.

462. *Myrtus communis* V. *Styr. ligur.*
in sylv. aprie. Nicaeni. maxime vulgaris. hujus Variet. Circa
Montalbani. Nicaeni. crescit.

465. *Punica granatum*
Frequent in Com. Nicaeni.

466. *Myrtus azarolus*
(*Myrtus* fol. obtus. subtrifid. dentatis.)
Comit. Nicaeni. indigena planta est.

467. *Orchis arachnites* V. *pl. ligur.*
in Coll. Calidior. Nicaeni.

468. *Epipactis atrorubens*
l. *aphylla* flor. inermi labello bicorni
in sylv. umbrat. Com. Nicaeni. Circa Sorpello.

469. ——— *Vulva*
l. *caul. pauciflor.* label. lanceolat. lineat. lineis undulatis.
in sylv. umbrat. montium Com. Nicaeni. minime rara.
Nativitatem circa Bivini.

470. *Pancratium maritimum* V. *pl. ligur.*
ad ripas prope litus maris ubique frequens.

471. *Narcissus tazetta*
N. *spatha multiflora* nectar. campanul. truncat. brevior
petalis foliis planis.
Nicaenium Collium indigena.

472. *Allium rotundum*.
A. *caul. planifol.* umbellifero stamin. brevissimis um-
bella subglobosa flor. lateral. nutant.
in Com. Nicaeni.

473. ——— *oreum* V. *pl. ligur.*
in olivetis & arvis, Com. Nicaeni. rarum.

474. ——— *salicifol.*
A. *caul. subterreus* fol. umbellifero flor. pendul. truncat.
stamin. simpl. coroll. aequantibus.
in Colle. Albani prope Nicaeni.

75. *Lilium pomponium*.
fol. spars. tubulatis, flor. reflexis, corol. revolut.

462. Myrtus communis V. Steg. Lign. in hyl. aprie. Micaleu. maxime vulgaris. hujus Vallet. Circa
in hyl. aprie.

46

41

4

41

4t

4

4

47

47

47

lilium
l. foli
in l

Jicla

Ap
Calyc
petalij
Hydrun
in po
Rad. re
Dial. g

Arpa

lythru

Ane

83. Ca

84. Lyp

Ha. S. W
milan
perv

475. *Lilium pomponium*.
l. fol. spars. tubulatis, flor. reflexis, corol. revolut.
in Comit. Nicæens. frequent.

476. *Scilla maritima*
ad maris litus arenosa & rupes inhabitat.

477. — *italica*
l. corymb. conferto subconico.
loc. lapidos. non macilentis & umbratis Comit. Nicæens.

478. *Aphyllanthes monspeliensis*.
Calycis etc squamis imbricatis compositis tubulosis corolla caryophyllæ
petalis intubum superius limbo patente instructum dispositis stigma
bifidum capsula turbinata 3-angulatis 3-locularis sem. ovata.
in pascuis siccis agri Nicæensis.
Rad. repens culm. pl. 2-palm. simplici cinct. basi vaginis utriusque foliar
dical. graminea flor. terminal solitariæ. curvæ.

479. *Asparagus acutifolius* V. pl. ligur.
uberrime in Comit. Nicæens.

480. *Lythrum hyssopifolia*.
l. fol. altern. linear. flor. hercand.
in tabulis hum. torventis paillois frequentissima.

481. — *thymifolia*.
l. fol. altern. linear. flor. ferrapetalis
in tabulis hum. paillois.

482. *Anemone hortensis*
et fol. digitalis
Nicæa occurrat.

483. *Cactus* ~~l.~~ *opuntia*.
ad rupes circa Solpello.

484. *Ephedra distachya*.
l. pedunc. opposit. ament. gemin.
Habitus fruticosus Caul. 3-ped. Vam. fol. equiseti fol. & vami si-
milares sub spicis stipul. 2. sicca lanceol. Vagina cylindrica.
perforata biconvex.
in Agro Nicæensi.

485. *Pinus pinea*
p. fol. geminis primordial. solitar. ciliat.
 in mont. Comit. Nicaeni.
486. *Juniperus oxycedrus* V. *p. ligur.*
 ubique in collibus & sylvis appici Nicaeni.
487. ——— *Lycia*
j. fol. tern. undique imbricatis ovatis obtusis.
 inter s. Dalmanzo & Soppello ad vapes.
488. ——— *phoenicea* V. *p. ligur.*
 frequens in Comit. Nicaeni.
489. *Quercus ilex.*
Q. fol. ovata. oblong. indivis. serratisque subt. incan. cortice integro.
 Sylvarum agri Nicaeni indigena.
490. ——— *uber*
Q. fol. ovata. oblong. indivis. serrata. subt. tomentosa. cortice nigro fungoso.
 in montanis Nicaeni habitat.
491. ——— *coccifera*
Q. fol. ovata. indivis. spinosa. dentata. glabra.
 in sylv. Nicaeni. variorum quam praecedens.
492. *Pistacia terebinthus*
p. fol. impar. pinnatis. foliol. ovata. lanceolata.
 in agro & collibus Nicaenibus ubique.
Foliot. 3juga inbreitalia dura fl. axill. Corymbos. peduncul. ramis. Vascis alterni fol. breviores.
493. ——— *lentiscus*
p. fol. abrupte pinnatis foliol. lanceolata.
 in aridioribus et saxis. Coll. Nicaeni. prope Mare.
Frutice parvus foliol. saepius 3. 6. juga varia integerr. perfoliis marginatis fl. axillar. Vascis amentis brevibus geminata sessilia dura divergentia. Fructus Nicaeni magnitud. primum rubelli dein nigri.
494. *Salicornia herbacea.* Calyce integer ventricos. Stam. 1. Sem. 1.
 ad maris Nicaeni (litus).
Frutice saepe cubito altior erectus. Articuli cylindrici apice capsulae spica articulata. Floretes terni approximati parvi lutei.
495. *Beta maritima*
B. flor. geminis in agro Nicaeni.

...
...
...
...
...

...
...
...
...
...

...
...
...
...
...

...
...
...
...
...
...
...
...
...
...

...
...
...
...
...
...
...
...

...
...
...
...
...
...
...
...
...
...

...
...
...
...
...

[Faint handwritten text, likely bleed-through from the reverse side of the page. The text is mostly illegible due to fading.]

B. flor. gemini in agro d. ca. ent.

[Marginal notes on the right side of the page, including botanical references and classifications.]
 Chenopodium
 in
 m.
 C. fol.
 in a
 m.
 folio
 in lo
 Alviptle
 Hemaphro
 m.
 Fol. al. 2 p.
 A. cau
 in Com
 aut erect sub
 par. sessile fr
 C.
 o
 Frutic. rham
 terminales rac
 Lo
 A.
 eli
 Plant. tota ter
 vult. virgat. f
 hemaphrod.
 compertus 5.
 Urtica pi
 l. fol. oppos
 comitat

496. *Chenopodium Scoparia*.
C. fol. linear-lanceol. plan. integerrimis.
in Com. Nicæni. frequent.

497. ——— *maritimum*.
C. fol. subulatis semi cylindricis.
in agro Nicæni.

498. ——— *fruticosum*
Salsola erecta fruticosa fol. filiform. obtusiuscul.
in solo argillaceo ad mare libet & secus flumen de Vau.

499. *Atriplex rosea*
Hermaphrod. Cal. 5-phyll. corol. 0. Stam. 5. Styl. 2-part. Sem. 1.
depress.
Fam. Cal. 2-phyll. coroll. 0. Stam. 0. Styl. 2-part. Sem. 1. compress. orbicul.
A. caule herbaceo fol. incanis serratis fruct. 4 angular. dentat.
in Com. Nicæni frequent.
Caul. erect. sub 2-ped. ramoliss. ram. diffus. albus, flor. axillares glo-
mer. ferriles fruct. compress. roseus.

500. ——— *Halimus*
A. caule fruticoso fol. dentatoidibus integris.
ad litus Nicæni.
Fruticosa v. hamis simil. ramos. fol. inordinata caruosa alba flores
feminales racemosi parvi purpureascent.

501. ——— *laciniata*.
A. caul. herbac. fol. dentatoid. dentat. subt. argenteis.
Etiam hæc agri Nicæni indigena planta est.
Planta tota tecta quasi epidermide seredente Caul. erect. teres
nudus virgat. flavescens fol. except. ultim. alterna spicæ terminal.
hermaphrod. anth. rubris fl. femin. axillares gemini Cal. fructus
compressus 5-dent. dente intermedio majore.

502. *Urtica pilulifera*.
U. fol. opposit. ovat. serrat. ament. fructifer. globos.
Comitatum Nicæni incolit.

503. *Rumex maritimus*
R. fl. hermaphrod. valvul. dentat. granifer. fol. lineari.
 in agro Nicaeni.
504. — *bucephalosphorus*
R. fl. hermaphrod. valvul. dentat. nudis pedicell. plan. reflex.
 in arvis et sabulosis in agro Nicaeni.
505. — *tuberosus*
R. fl. dioic. fol. lanceolato oblong. sagittat. hamis
patentibus radice tuberosa.
 in pratis agri Nicaeni.
506. — *multifidus*
R. fl. dioic. fol. hastat. articul. hastat.
 Circa Nicaenam & Villafrauca ad vupes.
507. *Polygonum maritimum.*
sp. fl. octand. hyginis axillaribus fol. oval. lanceolat.
semperviventibus caule protercente
 in arvis & sabulosis agri Nicaeni.
508. *Mecebrum paronychia*
Cal. 5. phyllis cartilagin. Stam. 5. Styl. 1. capsula 1. sperma.
3 fl. bract. nigr. argenteis obvallatis caul. procumb. fol. opposit.
 in sterilibus Com. Nicaeni.
509. — *capitatum.*
3 fl. bract. nigr. occultant. capitul. terminali. caul.
erectum. subt. villos.
 in agro Nicaeni.
510. *Camphorosma monspeliaca*
Cal. 4. fid. legum. altern. major. reflex. Stam. 4. Styl. bifid. capsul. 1. sperma.
C. fol. hirsutis linearibus.
 In arenosis & aridis agri Nicaeni praeterhumiditate Villafrauca
 & la Cantera
 Caul. pedat. fruticos. subvillos. albesc. fol. altern. serrit. odor. Champhor.
511. *Glauc Maritima.* *Cal. 5. fid. colorat. Stam. 5. Styl. 1. capsula*
1. locul. 5. valv. 5. sperma.
Rad. geunculata caul. erect. digital. ram. oppositis fol. serrit. linear-
lanceol. obtusiusc. integerr. subcarnea lacte v. v. dia glabra inter. opposit.
floral. altern. fl. axill. serrit. solitar. etc. alb. d. rub. purpure.

2. *Salvia Kali* V. fl. ligur.

50 Rumex maritimus 1 dentat. granifer. fol. line-v.

50

5.

50

50

50

50

51

5.

512. *Salsola Kali* V. fl. ligur.
Circa Nicæam.

513. ——— *Soda*.
J. herbac. patula fol. inermibus
in agro Nicæensi.

514. *Cithnus hypocithi*.
Monogyna Cal. tubul-campanul. 4. fid. colorat. super. Anther.
16. fert. Bacca 8 locul. polysperma.
Paralithica Cithi habitus otobanche
Caul. 3. 4 pollic. caudat. purpur. aut sublut. fol. aut squam
imbricatis sessilibus oblongis. Flor. termin. foliol. color. calyc.
mali punicea loricat. lutei flor. subiacent. squam. 7. opposit.
in densis purpurascens.
Circa Berra in Com. Nicæensi et Circa Nicæam lica
arbores et Cithor. radic. inracent.

515. *Juncus acutus* V. fl. ligur.
in locis maritimis et viciis.

516. *Theligonum Cynocrambe Monoclea Heptandra*
ad muror & sepes Nicæa.
Caul. diffusi flexuosi teretes succulenti summi arcuati opp.
positi divaricati fol. oleracea petiolata inter. oppos. super.
alterna ovata obtusiuscula nervosa sublineata loricata mar.
gine loricata. stipul. breves 3. dentatae latae membranac.
sem. volum.

517. *Ceratonia Siliqua* Vern. *Cavro* be
fruct. siliqua magna compressa pulpa repleta
ubique circa Nicæam Arbor
Folia alterna abrupt. pinnat. foliola bijuga subrecta sub.
orbiculata integerr. crassa firma. fl. axillar. racemosi sex.
lites.

518. *Zannichellia pumila*
in Com. Nicæensi ad rivulos.
Caul. tenues ramosi. crispiter densiorefformant fol. sessil. simpl.
capillar. integerr. ima alterna summa opposita vagina lata
orientem ramum excipit fl. axillar. solitar. sessil. caput.
subarbores.

519. *Ruppia maritima*

R. foliis acutissimis gramineis longior. fruct. fere umbell.
ad lit. Mar. Nicæus.
Caul. demer. herbac. articul. fol. simplic. altern. sessil li-
nearia flor. axillar. solitar. pedunculati.

520. *Xosteva Oceanica.*

? pericarpis, pedicellatis oliviformibus.
ad maris, nicæensis litus
Differt a sequenti radic. in pilum complicatis &
fruct. pedunculatis.

521. ——— *marina*

? pericarpis sessilibus.
ad litus maris Nicæensis.

Planta aquis marinis innata, flexuosa radice fasciculos folior.
angustorum emittit. fascicul. caulini, culmi graminei ad instar
singuli & singulis radicalibus ramulis prodeunt, valde infirmi
terre geniculati flexiles natantes. Fol. altern. vaginant li-
nearia verticaliter obtusa glabra mucosa sat ure viridia.
in apice ramulorum 2-3-4. folia unica vagina inclusa
radices semiuunciales linea paullo crassiores.

522. *Blitum Capitatum Monandra Monogynia*

Caul. 3-fidus corol. o. sem. 1. Calyc. - baccato fl. plur. in glomer.
colliguntur stabilius Chenopodii

fl. capitellis spicatis terminalibus

Non infrequens in Comit. Nicæens. sicut alia.

Caul. pedat. simplic. fol. longe petiol. hastata semibiloba quan-
doque integra dentata. globul. florales sessiles corcinei.

523. *Smilax aspera.* Vern. *Gramon d' Montagna*

J. caule aculeato fol. dentat. aculeatis cordatis 9-nerviis.

Vulgaris est in Comitatu Nicæensi.

Caul. scandens cirrhis ad bat. petiol. fol. altern. brev. petiol. sicc.
virgida acuta quandoque auriculata. Fl. axillar. glomerato-
spicati corol. alba odora baccae rubra vel nigra.

524. *Ruscus hypoglossum*

in Jarois, asperis incultis sterilibus Nicææ Lobelius.

5. *Olyvil alba.*

in Comitatu Nicaeni in vallis ad Villam Francam de Nica

51 *Amphibia maritima*

5.

5

5

5

52

Olyvil
in Com
pudol
Frabre
1. per
laten
3. Nivun
Arum
Radice
palmat
Alar
Phleu
p. spic
in A
Phalar
V
p. spic
p. spic
Alape
Pha
2.2
C.

525. *Olyvil alba*.
in Comitatu Nicaeni praesentem ad Villa Franca depre-
munt vulgari.

Frutice. 2. 3. pedalis striatus fol. altern. serris. acut. integerr.
fl. terminal. conferti subracemosi pedunculati calyce viridi
luteo fruct. esclaeo umbilicatus ruber.

526. *Nicum album* V. pl. ligur.
locis maritimis frequentior quam in predemontio.

527. *Arum avilatum*. V. pl. ligur.
in sylvis Nicaenibus copiose.
Radice globosa verrucosa fol. atro virentia rugosa petiol.
palmar. maculati flor pallide atro purpur.

528. *Nardus aristata* V. pl. ligur.
in agro Nicaeni

529. *Phleum Schoenoides*
ph. spic. ovalis obsoletis fol. breviss. mucronat. impleticant.
in Agro Nicaeni

530. *Phalaris Canavensis*
ph. panicula subovata spic. brevi glumis Canavensis
in Comit. Nicaeni.

531. — *paradosca*
ph. spica cylindrica foliulis mucronatis neutris plurimis infimis
proximis.
in Comit. Nicaeni.

532. *Alopecurus moissetianus*.
A. panicula pubescentia calyc. scabr. coroll. aristatis
Ad limit. agrorum in Comit. Nicaeni.

533. — *ciliatus*
A. culm. spicab. erect. glumis ciliatis.
C. Bellardi: in pratis editionibus di. Breglio.

534. — *litoreus*
phalar. calyce divaricato plano ovato ciliato glum. floral. interior.
2. dentata.
C. Bellardi: ad litora maris a clivis ad flum. le Var.

535. *Milium leudigerum*.

Ch. panicul. subspicat. floribus aristatis.
in Comit. Nicaeni.

536.

— *paradosicum*

Ch. flor. panicul. aristat.

in agro Nicaeni: Cl. Bellardi: ad ruppem mari proximam.

537.

Agrostis miliacea

A. spet. arboribus arist. femin. Vecta Nicaeni.

Circa Nicaenam legit Cl. Bellardi.

538

— *pinguis*

A. panicula compacta multica foliis involutis rigidis pan-
gibus superior. oblique oppositis.

Legit Cl. Bellardi locis aridis Maris Nicaeni: proxime.

539.

— *aculeata*

Stemul. culmi teretis ramo capitul. terminal. involu-
to brevissimo, rigido patente.
in agro Nicaeni.

540.

— *minima*

A. panicula multica foliis
in agro Nicaeni.

541.

Rotbolla incurva V. pl. Ligur.

Nicaea in agro mari proximo.

542.

Lagurus ovatus.

Gen. Calycis glumae aristatae aristae villosae. Folliculus bivalvis
valvula majori terminata duplici aristae cum tercia ad dorsum ac-
cedente valvula altera minori acuminata Semen oblongum folli-
culo arcte inclusum aristatum.

h. spica ovata aristata

in agro Nicaeni: frequens.

543.

— *cylindrica*

h. spica cylindrica multica

etiam agri Nicaeni: indigena planta est.

544.

Cynurus cuneus.

C. panicul. spic. sterilibus pendulis flor. aristatis.
in agro Nicaeni. loc. steril. mari proximo.

45. *Lolium tenuis*
L. spica multiflora peretis spiculis trifloris

5. *Milium leudigerum.*

6

52

51

53

54

51

5

5.

5

Montana
L. sp.
in

San. rigida
ad

Virgo Vir
B.
in

— m
B.
pro

— m
B.
in

Stomus
B.
in

—
h

—
C

— m
B.
gel
lo

—
B.
in

Festuca
Traceno
maculosa
in

45. *Lolium tenuis*
L. spica mutica teret. spiculis trifloris.
 in Com. Nicæensi, pratis & dumetis fasciatis.
546. *Poa rigida* V. M. Lign.
 ad limites agrorum secus vias locis aridis in Com. Nicæ. abunde.
547. *Phlox vivens*
Ph. spiculis ovalis calyce florulis (7) aequali
 in agro Nicæensi.
548. ——— *maxima.*
Ph. spiculis cordatis florulis 17.
 frequent in agris Comit. Nicæensi.
549. ——— *monispermatana.*
Ph. spica nutante simplic. spicul. altern. pedunculatis
subulatis ovalis calyce 5. flor.
 in agro Nicæensi.
550. *Bromus rubens.*
B. panic. fascicul. spicul. suberect. villis aristis erectis.
 in agro Nicæensi.
551. ——— *inermis*
B. panicula erecta spicul. suberectis, tubulatis nudis
inermis.
 (C. Bellardi circa Torpello).
552. ——— *Plukenetii*
B. fol. convolut. pungent. locust. binis terminalibus.
 C. Bellardi legit locis aridis Nicæa circa Villafanca.
553. ——— *madritensis*
B. panicul. varior. patulo-erect. spicul. linear. internod.
geminis pedicellis superne incrassatis.
 locis siccis circa Tashien in Com. Nicæensi.
554. ——— *Distachyos.*
B. spicis duabus erectis alternis.
 in Com. Nicæensi. ad vias & agrorum limites.
555. *Festuca pharicoides.*
F. racemo indiviso spicul. altern. suberect. teret. fol. involut.
mucronat. pungent.
 in pratis & ad sepes in Com. Nicæensi.

566. *Triticum junceum*.

T. calyc. truncat. 3. flv. fol. involutis.
ad Vias & loca sicca in agro Nicaeensi.

567. ——— *tenellum*

T. calyc. subq. flv. florent. mult. acut. fol. setaceis.
in Comitatu Nicaeensi.

568. ——— *unilaterale.*

T. calyc. unilateral. alternat. mult.?
Nascitur ad litus maris Nicaeensis.

569. ~~*Phymosus*~~ *avenarius.*

P. spica erecta arcta calyc. tomentis. florent. longioribus.
in arena ad maris litus.

560. *Hordeum geniculatum*

H. culm. decumbent. foliis oblecto summo spatheo
spica subjecto supremo nodo incurvato.
C. Bellardi legit ad litus maris Nicaeensis prope
portum di Limpia.

561. ——— *secalinum.*

H. florent. lateral. mascul. aristatis involucri setaceis
scabris
Habitat in agro Nicaeensi.

562. *Andropogon distachyon*

A. spic. 2. terminal. culmo indiviso
Ubique ad vasa et in lacubus Comit. Nicaeensis.

563. *Aegilops ovata.*

Gen. locusta 3. vel 4. flv. flores perfecti laterales & masculi in
terminis continens calyc. biglumis glum. ovata aristat. foliis
2. valv. valvula exterior ovata triplici aristata terminata interior
multica sem. oblongum valvulae foliis interiori aequale.
A. spica ovata calyc. omnibus 3. aristatis.
Loci siccis ubique in Comit. Nicaeensi.

564. ——— *vincialis.*

A. spica aristata calyc. inferioribus biaristatis.
in agro Nicaeensi.

in foliis

foliis

namque

ta

etiam

volucrum

officium

et in malis

foliis

minuta in

acervum

in foliis

[Faint, illegible handwriting throughout the page, likely bleed-through from the reverse side.]



565. *Cenchrus capitatus*
C. spica ovata simpliciter
in Monte Pyrae & circa Ioppello C. Bellardi.
566. *Cornu lachryma jobi*
in agro Nicaeni legit D. Giudice.
567. *Cavendishia*
C. caule foliis, capitulisq. hirsutis.
in agro Nicaeni foliis sylvaticis vulgaris.
568. *Scirpus mucronatus*
S. culmo 3. angulo modo acuminato spicis conglomeratis
sessilibus lateralibus.
Mareis in Maritimis Nicae.
569. — *maritimus*
S. culmo trigonato panicula conglomerata foliacea spicularum
ignavis 3. fidi intermedia subulata.
in com. Nicaeni frequens.
570. *Schannus mucronatus* V. pl. Lignv.
ad litora maris a Monaco ad Menton.
571. *Lichen Rocella*
L. fruticulosus solidus asphyllus 16. ramorum tuberculis alternis
ad Rupes maris Nicaeni.
572. *Fucus serratus* V. pl. Lignv.
in Mari Nicaeni.
573. — *Utricularia* V. utriusque
ibidem.
574. — *canaliculatus*
F. proinde plana dichotoma integerrima canaliculata
linearibus multiplicationibus tuberculatis in partibus obtusis
ad Rupes litoris Nicaeni.
575. — *nodosus* V. pl. Lignv.
ad mare Nicaeni.

576. *Fucus siliquosus*
F. frond. compressa ramosa fol. distich. alternis integerrimis fructi-
ficationibus pedunculatis oblongis mucronatis.
ad rupes littoris Nicæensis.
577. — *Delaginoideus* V. *sp. ligu.*
ad littus Nicæense
578. — *fœniculaceus* V. *sp. ligu.*
in rupibus littoris Nicæensis.
579. — *abrotanoideus* V. *sp. ligu.*
in mari Nicæensi
580. — *natus*
F. caud. filiformi ramosa fol. lanceol. serratis fructificat. glo-
bos. pedunculatis.
in mari Nicæensi
581. — *acinarius* V. *sp. ligu.*
in mari Nicæensi
582. — *linearis*
F. dichotomus planus linearis acutus vertic. ovatis spinis
Ad rupes maris Nicæensis.
583. — *fastigiatus*
F. frond. filiformi dichotoma ramosisima fastigiata
obtus.
ad rupes maris Nicæensis.
584. — *robustus* V. *sp. ligu.*
in mari Nicæensi.
585. — *ceranoides*
F. frond. plana dichot. integer. punct. lanceol. fructific.
hiberculat. bifid. terminal.
in mari Nicæensi.
586. — *dentatus*
F. frond. membranac. nerv. alternatim pinnatis. lacin. apice
evolutis
in mari Nicæensi.

576. *Ficus siligiosa*
T. fronde compressa, mox fol. distich. altern. integerrimis, vix

57

57

57

58

58

58

58

584

585

58

587. *Tucus splendens* *pleurocervoides*
F. parvus caulis. teret. humilitat. membranac. dilat. & lacert.
Ad ruper maris Nicaeensis.

588. — *ramosissimus*
F. caul. teret. ramosissimo fol. altern. subulat. veteicul.
volund. ianovic. sens. lib. & ciliatib.
Ad ruper maris Nicaeensis.

589. — *mucoides* N. Pl. lignu.
in mari Nicaeensi.

590. — *helminthochortus*
Comperva helminthoch. Dichotoma uncinat. & percaur.
ramis horizontalib.
Lechus est in Ruppib. maris Nicaeensis.

591. — *verrucosus*
F. filiformis teret ramoru. ramis simplicibus verticatis.
volundit lateralib.
ad ruper maris Nicaeensis.

592. — *longissimus*
F. teret. rubens minus ramor. in longum proterus
in mari Nicaeensi.

593. — *purpureus*
F. filiform. teret ramorill. ramis alternis ramul.
confertis brachiferis brachib. globosis.
ad mare Nicaense.

594. — *plicatus*
F. capillaris uniform. ramosissimus implicatus sub
diaphanum
ad ruper maris Nicaeensis.

595. — *corneus*
F. cartilagineus caule teret. intusculo compresso ramosi-
simo ramis subsplanatis laciniis acutis brachiferis
ad ruper maris Nicaeensis.

596. *Fucus confervoides* V. pl. Liguor.
in mari Nicaeni.

597. — *plumosus* V. pl. Ligu.
ibidem.

598. — *pinnatifidus*
F. frond. plan. vumol. vumol. dentato pinnatifidi
marginibus callosis
ad super mare Nicaeni.

599. — *cartilagineus*.
F. fronde cartilaginea compressa supradecomposita
pinnata laciniis linearibus
Ad super mare Nicaeni.

600. — *volubilis*
F. fronde plana spirali pectfol. repand. dentata.
ad mare Nicaeni.

601. — *spiralis*
F. frond. plan. dichot. integerr. punctata inferne
lineariter canaliculata vachisicant. tuberculat. geminis.
ad mare Nicaeni.

602. — *palmatus*
F. fronde palmata plana.
ad super mare Nicaeni.

603. *Ulva pavonia*
U. plana veniformis sessilis decussatim striata
in mari Nicaeni.

604. — *umbilicalis*
U. plana orbiculari sessili peltata coriacea
Habitat ad super mare.

605. — *intestinalis* V. pl. Liguor.
in mari Nicaeni.

606. *Alva compressa* V. pl. fig.

penal

supra

repand

chala infera
scula regenda

stadium / miala

la toria

ad vassellat hnt officium

5 10 15 20 25 30 35 40 45 50 55 60 65 70 75 80 85 90 95 100

60

60

60

60

60

60

606. *Ulva compressa* V. pl. lig.
Habitat in mari Nicaea.

607. — *Lactuca* v. ut lapon
ibidem.

608. — *linga*
ibidem.

609. *Conferva fasciculacea*.
Gen. Tubercula inaequalia in fibris capillaribus longissimis.
C. filament. aequal. ramis. ramis. ramulis. longiss.
Habitat in rupibus maris Nicaeensis.

610. — *tomentosa*
C. filament. aequal. ramis. ramis. ramulis. confert. fuscis.
ad litus maris Nicaeensis.

611. — *litorealis*
C. filam. aequal. ramis. elongat. scabris. caulis.
Habitat in rupibus litus Nicaeensis.

612. — *aeugina*
Plantae marinis litus Nicaeensis innascitur.
C. filament. ramis. mol. digit. brevior. viridiss.

613. — *Scoparia*
C. fil. proliferis fasciatis liliis
Habitat in mari Nicaeensi.

614. — *corallina*
C. filis geniculatis dichotomis
in litore maris Nicaeensis.

615. — *polymorpha*
C. filamentis geniculatis ramis fasciculatis
ad litus maris Nicaeensis.

616. — *rupestris*
C. filamentis geniculatis ramis. virid.
ad rupes litus Nicaeensis.

617. *Conserua elongata*
C. filam. genicul. ramisiss. ramuli longiss. distant. acut.
Habitat in litore Nicaeensi.

618. ——— *furca*
C. filam. geniculatiss. ramisissimiss. ramul. altern.
simplicibus
in litore Nicaeensi.

619. *Aristolochia rotunda*
A. fol. cordat. subellipt. obtus. caul. inferum. flor. solitar.
in sylvis collinis umbratis Nicaeensibus.

620. ——— *longa*
A. fol. cordat. petiol. integerr. obtusiuscul. caul. inferum
fl. solitariis
Sponte nascitur in vineis Nicaeensibus.

621. *Chamaerops humilis*
Ch. non palmat. plicat. stipit. spinos.
Palmam humiliss.
in agro Nicaeensi.
Auctantii

622. *Valeriana cornucopiae*
V. flor. grand. ringent. fol. ovat. sessil.
Nascitur in Comit. Nicaeensi.

623. *Antirrhinum origanifolium*. V. pl. Tende
supra l' Escavene.

624. *Scabiosa hybrida*
S. corollat. 4fid. radiant. fol. infer. pinnatis pinnul. obtus. dent. ramis integris
in arvis et pascuis agris Solypitellensibus.

625. *Senecio squallidus*
S. coroll. radiant. calyc. longior. integ. fol. pinnatis f. lacin. linear. distantibus.
Collis Nicaeensi. inhabit. et frequens est in Olivetis & locis lapideis
praesertim prope Saorgio.

23 Oct 1880

mat. 1880

mat. 1880

mat. 1880

mat. 1880

mat. 1880

mat. 1880

[Faint, illegible handwritten text, likely bleed-through from the reverse side of the page.]

600 *Centropomus viridis*
A large, deep-bodied fish, found in the
lagoon and near the shore.

601 *Sphyrna tiburo*
A large, deep-bodied fish, found in the
lagoon and near the shore.

602 *Scorpaenopsis diabolus*
A large, deep-bodied fish, found in the
lagoon and near the shore.

603 *Scorpaenopsis diabolus*
A large, deep-bodied fish, found in the
lagoon and near the shore.

604 *Chromis argenteus*
A large, deep-bodied fish, found in the
lagoon and near the shore.

605 *Valamona*
A large, deep-bodied fish, found in the
lagoon and near the shore.

606 *Chromis argenteus*
A large, deep-bodied fish, found in the
lagoon and near the shore.

607 *Chromis argenteus*
A large, deep-bodied fish, found in the
lagoon and near the shore.

608 *Chromis argenteus*
A large, deep-bodied fish, found in the
lagoon and near the shore.

627. *Hieracium prostratum*.
in prat. editioribus di. Soppello.
628. *Berberis ciliata*
3 fol. lingulat. spatulat, acutis, ciliatis, fruct. corymbois.
Inter la Scavona & Nizza loci dactylis et in spatui ad ora
sylvarum.
629. — *linifolia*
3 herbacea fol. linear. integerr. caul. dextratis, caul. paniculat.
corymb. hemisphaericis.
In arenosis collibus agr. Nicaensis versus flumen de Var.
630. *Brassica oleracea*
A radice caulescente tereti carnosa.
Sponte nascitur inter Villafranca & Monaco in rup. bui mari
proximis.
631. *Cheiranthus incanus*
C. fol. lanceol. integerr. obtus. incanis, siliquis apice
truncatis compressis, caule suffruticoso
Inhabitat rupes mari proximas prope Villafranca
atque spontaneus est.
632. — *micropsidatus*
C. siliquis apice 3-dentatis, fol. lyvatis.
in Sabulosis ad rivum Nicaensis litus.
633. *Euphorbia Taurinensis*
in vineis Nicaensibus frequentissima.
634. — *Papaver dubium*
Vulgare in com. Nicaens, ibique variat. fl. albo.
635. *Medicago circinnata*
M. legum. reniform. marg. dentat. fol. pinnat.
in agro Villafrancae Nicaensi. prope mare versus Monaco
636. — *laciniata*
Frequent in arvis circa Soppello & Breglio.

637. *Hedysarum Cristagalli*.
H. fol. pinnat. Legum. 1. specim. aculeatis crist. lacini lanceol. denticul.
ad mare Nicaeni: latus in Sabulosi.

638 *Cachrys Moriloni*.
Gen. Umbella magna involucri universi & partial. polyphyllum
flores aequales petala integra angusta; semina oblonga gibba
crassa ad latus non explanata substantia crassa fungosa.
C. foliol. linear. erectis binis aut ternis punctibus semiovatis
laxibus utrinque robustatis
Supra Breglio loco dicto Mauciana

639 *Smyrniolum olusatrum*.
Non infrequent in Comit. Nicaeni.

640 *Bupleurum Gerardi*
Frequentissime occurrit in Comit. Nicaeni.

641. *Bubon macedonicum*.
G. Umbella magna involucri universi & partial. polyphyllum
polyphyllum petala oblonga sem. ovata striata villosa.
B. foliol. rhombico-ovatis crenatis umbell. numerosissimis.
in pascuis siccis prope Nicaeam.

642. *Geranium malacoides*.
Non modo circa Nicaeam videtur sed frequent est in Comitatu
Nicaeni.

643 *Lavatera punctata*
L. caul. herbac. fol. subrotund. 5. angulatis. deinde hastatis; peduncul.
1. floris axillar. flor. campanulatis.
in agro Nicaeni: frequent
caulis rubet & minutis punctis albis aspersis & pedatis.

644. *Ranunculus agrarius*
R. radic. fibros. ann. fol. hirsut. semi-3-lob. petiol. acute serratis
calyc. hirsut. retroflexis
in agro Collinis Comit. Nicaeni ut praesertim in quibus
aqua aliquandiu persistit.

Real. pen.

Real. pen.

Real. pen.

Real. pen.

Real. pen.

Real. pen.

Real. pen.

Real. pen.

Real. pen.

Real. pen.

Real. pen.

Real. pen.

637. *Hedysarum Cristagalli*.
H. fol. pinnat. Legum. 1. p. 1. m. aculeatis crist. lac. lanceol. denticul.
1. 1. 1. 2. 1. 6. 1. 1. 1.

6

6

6

6

645. *Statice speciosa*
Limonium fol. lanceol. mucronat. radicalibus caul. ancipit di-
chotomo stipulis simplicibus.
Occurrit circa Ventimiglia.

646. *Galanthus autumnalis*.
leucos. spathe multiflora stylo filiformi.
in rupibus prope Villafanca Nicæensis.

647. *Allium magicum*.
A. caul. planifol. umbellifer. ramulo bulbifero stamin.
simplicibus
Com. Nicæens. indigena planta est.

648. ——— *monspellanum*.
A. densa teretifolia umbell. fastigiata. petal. linear. fol. lanceol. stamin.
tubulatis
in Olivetis Villafanca Nicæens.

649. *Fritill. involucr.* N. sp. Monvey.
in Com. Nicæens. supra Broylio

650. *Atriplex hastata*.
A. caul. herbaceo fol. 3. angular. calyc. valvat. femineis magnis
deltoidibus sinuatis 3. angular. subaspreis
in arena litore Nicæensis.

651. *Aquilegia miliacea*
in Com. Nicæens. frequens est secus torrentes locis siccior.

652. *Stipa juncea*
St. arist. nud. rect. calyc. sem. longior. fol. intus (aristibus).
locis siccioribus calidior. in agro Nicæens. prope Villafanca.

653. ——— *aristella*.
St. aristis nudis rectis calyce duplo longioribus gemmin. lanatis.
Frequens locis rupestribus & calidioribus C. Nicæ. abunde circa
Villafanca.

654. *Brixa minor*
Frequentest secus arva locis calidioribus agri Nicæensis.
655. *Bromus plukenetii*
Frequentissimus est in collibus calidioribus & siccis Com. Nicæensis.
656. *Melica Benthini*
M. spica ramosa unilateralis fertilibus floribus petalo subciliato
provenit locis superstitibus aridis in Com. Nicæensis.
657. *Avena fatua*
Frequentissima est secus agros & in paucis collibus Calidiorum
Nicæensium.
658. — *bromoides*
A. subspicata spiculis binatis altera pedunculata aristata
varicatis calyc. 8-floris.
659. *Holcus arenarius*
H. spica conica tomentosa lucida glumis 6-floris hermaphrodito testili masculo pedunculata.
in arena ad mar. Nicæensis litus.
660. *Agropyron truncatis*
provenit præsertim secus agros inter Sospello & Castiglione
661. *Cyperus distachyos*
C. culmo triquetronudo spiculis sessilibus lateralibus
radice perenni.
locis spatiosis inter Nicam & le Var.
662. — *globosus*
C. rad. perenn. caespitosus planifol. culm. triquet. fol.
brevior. capit. globos. terminal. involver. byssillo.
in pratis spongiosis secus le Var.

654. *Arctostaphylos*

655.

656.

657

658

659

660

661

662

653. *Lichen*
h.
m.
in

654. *Arctostaphylos*

655. *Bulbo*
Cort. in
super.
in
in

656. *Tulipa*
T. fl. 23
in

657. *Anemone*
A. fol. v.
Mar.
Fl.

658. *Penstemon*
P. fl. 23
Calyc.
D. 23
in

659. *Geni*
G. fol.
Jaki

663. *Lichen diffusus*
L. pyxidatus endiv. (virg. foli. in fundibul. explanati
fructiferi
in pascuis sterilibus & dumetis Com. Nicæensis.
Appendicis.

664. *Anchusa officinalis*
A. fol. lanceol. spic. imbricatis secundis
Nascitur in Com. Nicæensi ad Vinas.

665. *Balbocodium Vernum.*
Corol. in fundibul. form. 6. petal. ungu. angust. staminifer. Corol.
super. Genus Colchico a quo differt petalis divisis usque
ad radicem ^{unguibus} petalis staminiferis.
in Com. Nicæensi reperit Marchia De-Suffren.

666. *Tulipa geneviana.*
T. fl. erecto fol. ovat. lanceolatis.
in collibus horisque incultis circa Nicæam.

667. *Anemone Coronaria*
A. fol. radical. ternato decompositis involucri folioso.
Nascitur in montibus Nicæa locis incultis.
Flor mire ludit color. Violac. subalbido rubro & stramineo.

668. *Jeneverium tomentosum*
J. fol. crenat. tomentos. flavicantibus floribus aureis
calycibus & foliis floralibus intente flavis distinguitur a
J. polio.
in Com. Nicæensi.

669. *Genista Scoparia*
G. fol. simplicibus hirsutis ramis to. angulatis cinereis
Sabi frequent in Com. Nicæensi. loc. montan. calidior.

670. *Tragopogon porrifolium*.

*T. calyc. coroll. radio sequi: longioribus fol. integris strictis pedunc.
superne incrassatis.*

Nascitur in Com. nicdensi Observ. D. Galli.

Flor. purpureo corat. prodit.

671. *Acer opulifolium*

A. fol. lobis subrotundis obtuse dentatis fructibus cymosis.

*in Com. Nicdensi. Sponte nascitur in rupibus a Fontana ad
Saurgium.*

672. *Pteris Cretica*

*pp. frond. pinnat. pinn. opposit. lanceol. serrat. bas. angustat. in pinn. sub-
tripartitis*

*in Vallecula umbrosa agri Nicdensi detescit. Cl. Desuffren.
etiam in sylvaticis umbratis Tenda.*

Gunei Die 11.
Junii 1800



Integritas

1800

Integritas

Integritas

Integritas

Integritas

Integritas

Integritas

Integritas

Integritas

Integritas

Integritas

Integritas

Integritas

Integritas

Integritas

Integritas

Integritas

Integritas

67

671

672



6

67

67








































Handwritten text in Arabic script, likely a library or archival label, located in the bottom right corner of the page.